

Diffuse leptomeningeal enhancement in neurosarcoidosis-related longitudinally extensive myelitis

Realce leptomenígeo difuso na mielite longitudinalmente extensa relacionada à neurosarcoidose

Fabiano Ferreira de Abrantes¹ Marianna Pinheiro Moraes de Moraes¹ José Luiz Pedroso¹
Orlando G. Barsottini¹

¹Universidade Federal de São Paulo, Escola Paulista de Medicina,
Departamento de Neurologia e Neurocirurgia, São Paulo SP, Brazil.

Address for correspondence Fabiano Ferreira de Abrantes
(email: fabianofabrantes@hotmail.com)

Arq. Neuropsiquiatr. 2023;81(11):1016–1017.

A 35-year-old man presented with a 1-month history of bilateral lower limb weakness and numbness. On examination, he had paraparesis with lower limbs hyperreflexia, and T5 sensory level. Magnetic resonance imaging (MRI) of the spine revealed a longitudinally extensive myelitis (LETM) with a diffuse leptomeningeal enhancement (**►Figure 1**). Thorax computed tomography (CT) disclosed bilateral hilar lymphadenopathy. A transbronchial lymph node fine-needle biopsy revealed noncaseating granulomas. Considering this finding, probable neurosarcoidosis was diagnosed.

Leptomeningeal involvement is a remarkable finding in neurosarcoidosis.¹ When central nervous system (CNS) involvement occurs concomitantly with a widespread leptomeningeal enhancement, sarcoidosis should be considered as a possible differential diagnosis.²

Authors' Contributions

FFA, MPMM: conceptualization, writing – original draft, writing – review & editing; JLP, OGB: conceptualization, writing – review & editing.

Conflict of Interest

The authors have no conflict of interest to declare.




Figure 1 T2-weighted (A) image of the cervical spine demonstrates longitudinally extensive myelitis (between the dashed lines). Contrast-enhanced T1-weighted (B) image with an extensive posterior spinal enhancement (arrow), and an anterior meningeal enhancement (arrowhead). Contrast-enhanced T1-weighted (C) of the thoracic spine disclosing an extensive posterior leptomeningeal enhancement (arrowheads), and (D) a circumferential involvement of the thoracic leptomeninges (arrowhead).

received
April 18, 2023
accepted
May 3, 2023

DOI [https://doi.org/
10.1055/s-0043-1772604.](https://doi.org/10.1055/s-0043-1772604)
ISSN 0004-282X.

© 2023. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution 4.0 International License, permitting copying and reproduction so long as the original work is given appropriate credit (<https://creativecommons.org/licenses/by/4.0/>).

Thieme Revinter Publicações Ltda., Rua do Matoso 170, Rio de Janeiro, RJ, CEP 20270-135, Brazil

References

- 1 Murphy OC, Salazar-Camelo A, Jimenez JA, et al. Clinical and MRI phenotypes of sarcoidosis-associated myelopathy. *Neurology(R) neuroimmunology & neuroinflammation.* NLM (Medline); 2020;7
- 2 Dutra LA, Braga-Neto P, Oliveira RA, Pedroso JL, Abrahão A, Barsottini OGP. Neurosarcoidosis: guidance for the general neurologist. *Arq Neuropsiquiatr* 2012;70(04):293–299. Doi: 10.1590/s0004-282x2012000400014