



Prolonged Fasting as a Cause of Deep Vein Thrombosis: A Case Report

Guillaume Roberge¹  Carine Samson² Grégoire Le Gal³  Anthony Calabrino¹

¹Centre d'Excellence des Maladies Vasculaires, Centre Hospitalier Universitaire de Québec, Université Laval, Hôpital Saint-François d'Assise, Québec, Canada

²MaClinique Médicale Lebourgneuf, Québec, Canada

³Department of Medicine, Ottawa Hospital Research Institute, University of Ottawa, Ottawa, Ontario, Canada

Address for correspondence Guillaume Roberge, MD, DRCPC, Centre d'Excellence des Maladies Vasculaires, Centre Hospitalier Universitaire de Québec, Université Laval, 10, rue de l'Espinay, Québec, Canada G1L 3L5 (e-mail: guillaume.roberge.2@ulaval.ca).

TH Open 2023;7:e94–e96.

Abstract

Background Intermittent fasting is becoming more popular as health benefits are described in recent literature. Various forms of fasting exist, one of them involving a zero-calorie diet and drinking only water. However, the safety of water-only fasting is still not well established. We report a case of a man who developed a lower limb deep vein thrombosis at the end of a 2-week water-only fasting and characterized by an initial period of 5 days of no food and no water intake. We reviewed literature regarding potential links between fasting and venous thromboembolism (VTE).

Clinical Approach We believe that fasting can induce important dehydration, leading to hypercoagulability and then contribute to the development of a venous thrombosis. The patient was treated with apixaban for 3 months as is recommended in patients with a provoked event caused by a transient risk factor. No thrombotic recurrence was observed during the 6-month follow-up.

Conclusion The public needs to be aware of the potential life-threatening complications associated with important dehydration in the setting of medically unsupervised fasting, and these might include VTE. Whether a VTE with dehydration as the only identified risk factor should be approached as a low recurrence risk situation or not still needs to be clarified.

Keywords

- ▶ fasting
- ▶ thrombosis
- ▶ venous thrombosis
- ▶ dehydration
- ▶ anticoagulation

Introduction

Intermittent fasting is becoming more popular as health benefits are described in recent literature. Weight loss, reduction in abdominal fat, decrease inflammation, lowering blood pressure, and reduction of insulin resistance are some of the observed effects of dietary restriction.¹ Various forms of fasting exist but medical supervision is usually recommended, especially when food restriction is more important or for a prolonged period of time.^{2–4}

Water-only fasting is defined by restriction of every type of food (zero-calorie diet) except water. In the 1960s and

1970s, fasting duration of 60 days and more was considered for important weight loss, but this practice has been abandoned due to important complications and deaths. At that time, fasts were improperly done due to lack of physicians' and researchers' knowledge about safety and risks.⁵ This kind of fast has also been described in nonobese persons for religious purposes.^{6,7}

Nevertheless, potential health benefits are still reported after shorter course of water-only fasting done under appropriate medical supervision.^{4,8,9} However, safety of such food restriction is still not well established and relies on limited data.^{4,5}

received
January 31, 2023
accepted after revision
March 21, 2023

DOI <https://doi.org/10.1055/s-0043-57225>.
ISSN 2512-9465.

© 2023. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution License, permitting unrestricted use, distribution, and reproduction so long as the original work is properly cited. (<https://creativecommons.org/licenses/by/4.0/>)
Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany

We describe the case of a man who developed a right lower limb proximal deep vein thrombosis (DVT) in the context of unsupervised water-only fasting.

Case Presentation

A 59-year-old male, with a body weight of 70 kg and body mass index of 24 kg/m², presented with a progressive right lower limb pain and edema starting at the end of a voluntary 2-week water-only fasting period. He was not known for any relevant past medical or family history. In the last year, the patient had undergone three medically unsupervised water-only fasting periods of 1 week for potential health benefits purposes. He had progressively increased not only the frequency of these fasts but also their duration up to 2 weeks. In his second 2-week fasting, the patient was still walking 5 km per day, lost around 10 kg (14% of total body weight), and started drinking 500 mL of water per day only on day 6 (i.e., no water for the first 5 days). He did not take any food supplements or weight loss drugs. The last 3 days of the fast (days 12 to 14), the patient reduced his activities and his walks because of fatigue but he was not bedridden. While he was resuming food intake (only fruit juice), he developed his right leg symptoms. Ten days after, he sought medical attention, and a diagnosis of right lower limb proximal DVT (peroneal veins up to the common femoral vein) was made. There were no identified risk factors for venous thromboembolism (VTE) (no recent surgery, trauma or admission to hospital, no long travel, no family history of VTE, no known comorbidities, no obesity, no active smoking, no varicose veins, no recent coronavirus disease infection) and blood work was normal at diagnosis. Thrombophilia testing was not done. Apixaban was initiated and the patient's symptoms progressively improved over the next weeks. At 3 months, anticoagulation was discontinued, and a repeated venous ultrasound revealed a residual thrombus without recanalization in the peroneal and femoral veins, but popliteal vein was partially recanalized.

After 6 months of follow-up, he was still presenting slight right leg edema, heaviness while walking, and new varicose veins, but did not have a venous thrombosis recurrence.

Discussion

Fasting Causes Dehydration

Complex physiologic processes occurring with fasting have been hypothetically suggested as a cause of observed increased diuresis and potential resulting dehydration.¹⁰ The decrease of glucose intake impairs sodium reabsorption as glucose-sodium cotransport in renal proximal tubules is inhibited. This ensued natriuresis is not compensated enough by the distal tubular sodium reuptake and then explains resulting osmotic diuresis and subsequent dehydration.¹⁰⁻¹²

As the patient was taking no calorie and 500 mL of water per day starting only on day 6, and experiencing a 10-kg weight loss in 14 days (weight that he regained in 1 week after restarting food intake), significant dehydration was most likely present. Also, it is improbable that only the first 5 days with no intake (no food, no water) explain the whole

dehydration as weight loss was progressive over the 14 days. Despite water intake, raised hematocrit and blood urea nitrogen were reported after water-only fasting in two cases, reflecting potential dehydration.^{7,13}

Dehydration is a Risk Factor of VTE

Dehydration is one of the potential explanations put forward to explain the increased VTE incidence observed after prolonged travel.¹⁴ This state likely results in hypercoagulability induced by hemoconcentration and hyperviscosity as reflected in part by increased biochemical parameters such as hematocrit, plasma proteins, plasma, and urine osmolality.¹⁵ Other conditions in which VTE likely occurs because of dehydration have been described such as following ischemic strokes, intense exercises, and gastroenteritis.¹⁶⁻²⁰ Also, according to some authors, seasonal variation of VTE incidence could hypothetically be related to dehydration, occurring more often with higher temperature.²¹

Some literature suggests that theoretically, medically supervised water-only fasting could in fact decrease thrombosis risk. Decreased platelet formation and activity was observed, but this was tested only after 7 days of fasting.⁹ Also, increased fibrinolytic activity was described in 12 healthy men after a 60-hour fast in which only water was allowed.²² However, vascular clinical outcomes under these circumstances were not reported. Of note, these studies do not apply to our unsupervised patient after a 2-week fast.

Important dehydration in this healthy and active patient most likely led to a state of hypercoagulability. It is unknown if an undiagnosed underlying predisposition to thrombosis such as a hereditary thrombophilia was present or if fasting can affect coagulability in another way. We believe that important dehydration in the context of fasting can represent on its own a transient risk factor for VTE. However, it is unknown if it should be interpreted as a major or minor one and deemed at low risk of recurrence.²³ We have not found any previous report of thrombosis complications following fasting.^{5,6}

Immobility could also represent a risk factor in fasting if the patient becomes confined to bed because of weakness. However, we highly doubt it explains the VTE in our case. Indeed, despite the reported reduction in his activities in the last 3 days of his fast, the patient was still mobile.

A concerted decision was made to treat 3 months with anticoagulation, as recommended with provoked VTE caused by a transient risk factor.²⁴ No recurrence was observed up to 3 months after stopping the treatment.

There are limitations to the present case, as exact hydration status was not established at the time the DVT occurred considering the patient had regained his usual weight at diagnosis. Also, we do not have any available blood test results at the end of the 2-week fast in correlation with dehydration. This is only one case and firm conclusions cannot be drawn.

Conclusion

While literature suggests potential benefits from intermittent fasting, the public needs to be aware of the potential life-threatening complications associated with important

dehydration in the setting of medically unsupervised fasting, and these might include VTE. Whether VTE caused in the context of dehydration as the only identified risk factor should be approached as a low recurrence risk situation or not still needs to be clarified.

Highlights

- Intermittent fasting is getting more popular.
- Medically unsupervised fasting can lead to significant dehydration.
- Dehydration is a risk factor for venous thromboembolism.
- Risk of thrombosis recurrence and best management in this setting is uncertain.

Conflict of Interest

G.R. has received honoraria from BMS, Pfizer, Bayer, Servier, Astra Zeneca, L'Académie, and Leo Pharma.

C.S. has no competing interest.

G.L.G. has received grants from Pfizer and Bristol-Myers Squibb and honoraria (not taken as salary) from Pfizer, Sanofi, and Aspen Pharma.

A.C. has received honoraria from L'Académie, BMS/Pfizer, Bayer, Boehringer-Ingelheim, CPD Network, Servier, and LEO Pharma.

References

- de Cabo R, Mattson MP. Effects of intermittent fasting on health, aging, and disease. *N Engl J Med* 2019;381(26):2541–2551
- Longo VD, Mattson MP. Fasting: molecular mechanisms and clinical applications. *Cell Metab* 2014;19(02):181–192
- Wilhelmi de Toledo F, Buchinger A, Burggrabe H, et al. Fasting therapy—an expert panel update of the 2002 Consensus Guidelines Revised Consensus Guidelines for Fasting Therapy 435. *Forsch Komplement Med* 2013;20:434–443
- Ogłodek E, Pilis Prof W. Is water-only fasting safe? *Glob Adv Health Med* 2021. Doi: 10:21649561211031178
- Finnell JS, Saul BC, Goldhamer AC, Myers TR. Is fasting safe? A chart review of adverse events during medically supervised, water-only fasting. *BMC Complement Altern Med* 2018;18(01):67
- Kerndt PR, Naughton JL, Driscoll CE, Loxterkamp DA. Fasting: the history, pathophysiology and complications. *West J Med* 1982; 137(05):379–399
- Brett AS, Nesbit RM. A 40-day water-only fast by a Pentecostal woman: clinical and religious observations. *Am J Med Sci* 2013; 345(05):418–420
- Goldhamer A, Lisle D, Parpia B, Anderson SV, Campbell TC. Medically supervised water-only fasting in the treatment of hypertension. *J Manipulative Physiol Ther* 2001;24(05): 335–339
- Fang Y, Gu Y, Zhao C, et al. Impact of supervised beego, a traditional Chinese water-only fasting, on thrombosis and haemostasis. *BMJ Nutr Prev Health* 2021;4(01):4–17
- Heyman SN, Bursztyn M, Szalat A, Muszkat M, Abassi Z. Fasting-induced natriuresis and SGLT: a new hypothesis for an old enigma. *Front Endocrinol (Lausanne)* 2020;11:217
- Maagøe H. Changes in blood volume during absolute fasting with and without sodium chloride administration. *Metabolism* 1968; 17(02):133–138
- Spark RF, Arky RA, Boulter PR, Saudek CD, O'Brian JT. Renin, aldosterone and glucagon in the natriuresis of fasting. *N Engl J Med* 1975;292(25):1335–1340
- Korbonits M, Blaine D, Elia M, Powell-Tuck J. Metabolic and hormonal changes during the refeeding period of prolonged fasting. *Eur J Endocrinol* 2007;157(02):157–166
- Gavish I, Brenner B. Air travel and the risk of thromboembolism. *Intern Emerg Med* 2011;6(02):113–116
- Chee YL, Watson HG. Air travel and thrombosis. *Br J Haematol* 2005;130(05):671–680
- Kelly J, Hunt BJ, Lewis RR, et al. Dehydration and venous thromboembolism after acute stroke. *QJM* 2004;97(05):293–296
- Hull CM, Harris JA. Cardiology patient page. Venous thromboembolism and marathon athletes. *Circulation* 2013;128(25): e469–e471
- Richard S, Lacour JC, Frotscher B, Enea A, Mione G, Ducrocq X. Report of a recurrent cerebral venous thrombosis in a young athlete. *BMC Neurol* 2014;14(01):182
- Lee KW, Lip GYH. Acute versus habitual exercise, thrombogenesis and exercise intensity. *Thromb Haemost* 2004;91(03):416–419
- Kuo EY, Chung MH, Lin HJ, Chen KT. Gastroenteritis followed by headache: a rare presentation of cerebral venous thrombosis. *J Acute Med* 2012;2(03):92–94
- Elias S, Hoffman R, Saharov G, Brenner B, Nadir Y. Dehydration as a possible cause of monthly variation in the incidence of venous thromboembolism. *Clin Appl Thromb Hemost* 2016;22(06): 569–574
- Miettinen M. Effect of fasting on fibrinolysis and blood coagulation. *Am J Cardiol* 1962;10(04):532–534
- Kearon C, Ageno W, Cannegieter SC, Cosmi B, Geersing G-J, Kyrle PASubcommittees on Control of Anticoagulation, and Predictive and Diagnostic Variables in Thrombotic Disease. Categorization of patients as having provoked or unprovoked venous thromboembolism: guidance from the SSC of ISTH. *J Thromb Haemost* 2016; 14(07):1480–1483
- Stevens SM, Woller SC, Kreuziger LB, et al. Antithrombotic therapy for VTE disease: second update of the CHEST Guideline and Expert Panel Report. *Chest* 2021;160(06):e545–e608