

Editorial

Ventilator-induced Diaphragmatic Dysfunction in Patients with Traumatic Brain Injury

Parmod K. Bithal¹ Girija P. Rath¹

¹Department of Neuroanaesthesiology and Critical Care, All India Institute of Medical Sciences (AIIMS), New Delhi, India

J Neuroanaesthesiol Crit Care 2023;10:147-148.

Moderate to severe traumatic brain injury (TBI) represents one of the most important causes of death and disability. Although morbidity and mortality in these patients are principally due to their primary pathology, non-neurological complications, including respiratory dysfunction, are frequent contributors. Therefore, all TBI patients with moderate and severe brain damage are managed with invasive mechanical ventilation. Notwithstanding the fact that mechanical ventilation to support respiration is a life-saving intervention in critically ill patients, weaning becomes progressively more difficult with the increasing duration of ventilation. Ventilator-induced diaphragmatic dysfunction (VIDD) is believed to be one of the major contributors to the weaning difficulties in intensive care unit (ICU) patients. It significantly influences the duration of mechanical ventilation, weaning failure, morbidity, and mortality. The combination of prolonged mechanical ventilation and the effects of extended immobility in the ICU causes significant changes to muscle fibers, reducing both respiratory and peripheral muscle strength.¹ Administration of muscle relaxants and/or steroids in mechanically ventilated patients further exaggerates the VIDD. The prevalence of diaphragm dysfunction has been reported to be twofold higher than the prevalence of ICU-acquired weakness.² Although VIDD has received considerable attention in critically ill patients in general ICU, it has not attracted sufficient diligence in the moderate to severe TBI patient population despite their requirement for prolonged mechanical ventilation.³

The diaphragm thickness decreases rapidly during the initial several days of mechanical ventilation in 40% of patients, and lower levels of inspiratory effort and higher levels of ventilatory support predict this decrease. Diaphragmatic thickness has been shown to reduce by 6 to 7.5% per day in mechanically ventilated patients.⁴ The diaphragm thickness negatively correlates with the ICU length of stay and positively correlates with the before/after rehabilitation functional scores and the change in functional independence measure scores.⁵ Diaphragmatic weakness and dysfunction decrease

Address for correspondence Girija Prasad Rath, MD, DM, Department of Neuroanaesthesiology and Critical Care, All India Institute of Medical Sciences (AIIMS), New Delhi, India (e-mail: girijarath@aiims.edu). DOI https://doi.org/ 10.1055/s-0044-1782223. ISSN 2348-0548. inspiratory pressure, resulting in difficult weaning.⁶ Patients with VIDD are significantly more likely to experience newonset sepsis and ventilator-associated pneumonia, are more likely to have extubation failure, and require a more prolonged duration of mechanical ventilation.⁷ Ultrasound imaging of the diaphragm is a reliable and reproducible tool for diagnosing neuromuscular diaphragm dysfunction.

Several studies have demonstrated that maintaining spontaneous respiration during mechanical ventilation can reduce VIDD. Under forced breathing, the increased inspiratory effort minimizes the incidence of diaphragmatic dysfunction to 11.5%, emphasizing that prolonged mechanical ventilation impairs diaphragmatic function independent of underlying lung disease.⁸

In brain-dead organ donors, a combination of 18 to 69 hours of complete diaphragm inactivity and mechanical ventilation resulted in marked atrophy of myofibers, which is consistent with increased diaphragmatic proteolysis. The diaphragm biopsy of these brain-dead donors showed marked atrophy of both slow- and fast-twitch fibers.⁹ Increased activity of the ubiquitin-proteasome pathway, marked decreases in myosin heavy chains (MyHCs), and atrophic AKT-FOXO signaling play important roles in eliciting the myofiber atrophy and decreases in diaphragm force generation associated with prolonged human diaphragm disuse.¹⁰ Mechanical ventilation also increases oxidative stress in the diaphragm, causing decreased protein synthesis and increased activity of proteolytic pathways, further accelerating the atrophy process.¹¹ Other comorbidities and metabolic stressors could negatively affect patients with VIDD as well.¹²

There is a need for alternative therapeutic strategies for preventing VIDD, including antioxidants and inotropic agents such as theophylline, digoxin, levosimendan, angiotensin receptor blockers, and phrenic nerve pacing.^{13,14} Amino acid intake during the first 24 hours of stay in the ICU may represent a significant, modifiable risk factor for VIDD; it may have a direct causal effect on mortality.⁷

(https://creativecommons.org/licenses/by/4.0/)

Thieme Medical and Scientific Publishers Pvt. Ltd., A-12, 2nd Floor, Sector 2, Noida-201301 UP, India

^{© 2024.} The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution License, permitting unrestricted use, distribution, and reproduction so long as the original work is properly cited.

In this issue of the journal, Das et al¹⁵ have published their observations on diaphragmatic function (VIDD and diaphragmatic thickness) in mechanically ventilated adult patients with moderate to severe TBI. While the study deserves appreciation, it has raised some pertinent issues. Contrary to the available literature on VIDD in general ICUs, the authors could not establish any significant correlation between VIDD and length of hospital stay. Is it due to the underlying pathology (TBI vs. non-TBI) having a role in the variable progression of VIDD and, thereby, length of hospital stay? The lack of a control group comprising TBI patients not subjected to mechanical ventilation or patients undergoing mechanical ventilation for reasons other than TBI limits the ability to isolate the specific contribution of mechanical ventilation to observed diaphragmatic changes. A relatively modest sample size of 40 patients, while providing initial insights, may lack statistical robustness. The study focuses on the initial week of ICU admission without extending the observation period to capture the peak of VIDD, which literature suggests occurs between the first and second weeks of mechanical ventilation. A more extended follow-up period would provide a more comprehensive understanding of the temporal progression of diaphragmatic dysfunction. Enrolling a larger sample size of moderate to severe TBI patients is essential to addressing this issue. Another area of research could be on automatic and spontaneous breathing-synchronized phrenic nerve stimulation (invasive and noninvasive) in combination with invasive or noninvasive ventilatory support forms.¹⁶ Future studies should also focus on nutrition and various modes of invasive mechanical ventilation vis-a-vis their impact on VIDD. It should also be studied whether gender has dissimilar courses of VIDD, given their distinct hormonal profiles.

Conflict of Interest None declared.

References

- 1 França EE, Ferrari F, Fernandes P, et al. Physical therapy in critically ill adult patients: recommendations from the Brazilian Association of Intensive Care Medicine Department of Physical Therapy. Rev Bras Ter Intensiva 2012;24(01):6–22
- 2 Dres M, Dubé BP, Mayaux J, et al. Coexistence and impact of limb muscle and diaphragm weakness at the time of liberation from

mechanical ventilation in medical intensive care unit patients. Am J Respir Crit Care Med 2017;195(01):57-66

- ³ Futier E, Constantin JM, Combaret L, et al. Pressure support ventilation attenuates ventilator-induced protein modifications in the diaphragm. Crit Care 2008;12(05):R116
- 4 Zambon M, Beccaria P, Matsuno J, et al. Mechanical ventilation and diaphragm atrophy in critically ill patients: an ultrasound study. Crit Care Med 2016;44(07):1347–1352
- 5 Guzel S, Umay E, Gundogdu I, Bahtiyarca ZT, Cankurtaran D. Effects of diaphragm thickness on rehabilitation outcomes in post-ICU patients with spinal cord and brain injury. Eur J Trauma Emerg Surg 2022;48(01):559–565
- 6 Powers SK, Wiggs MP, Sollanek KJ, Smuder AJ. Ventilator-induced diaphragm dysfunction: cause and effect. Am J Physiol Regul Integr Comp Physiol 2013;305(05):R464–R477
- 7 Pu H, Doig GS, Lv Yu, et al. Modifiable risk factors for ventilatorinduced diaphragmatic: a multicenter observational study. BMC Pulm Med 2023;23:343–352
- 8 Anzueto A, Peters JI, Tobin MJ, et al. Effects of prolonged controlled mechanical ventilation on diaphragmatic function in healthy adult baboons. Crit Care Med 1997;25(07):1187–1190
- 9 Levine S, Nguyen T, Taylor N, et al. Rapid disuse atrophy of diaphragm fibers in mechanically ventilated humans. N Engl J Med 2008;358(13):1327–1335
- 10 Levine S, Biswas C, Dierov J, et al. Increased proteolysis, myosin depletion, and atrophic AKT-FOXO signaling in human diaphragm disuse. Am J Respir Crit Care Med 2011;183(04):483–490
- 11 Shanely RA, Zergeroglu MA, Lennon SL, et al. Mechanical ventilation-induced diaphragmatic atrophy is associated with oxidative injury and increased proteolytic activity. Am J Respir Crit Care Med 2002;166(10):1369–1374
- 12 Demoule A, Jung B, Prodanovic H, et al. Diaphragm dysfunction on admission to the intensive care unit. Prevalence, risk factors, and prognostic impact: a prospective study. Am J Respir Crit Care Med 2013;188(02):213–219
- 13 Kim WY, Lim CM. Ventilatory induced diaphragmatic dysfunction: diagnosis and role of pharmacological agents. Respir Care 2017;62(11):1485–1491
- 14 Hall SE, Ahn B, Smuder AJ, et al. Comparative efficacy of angiotensin II type I receptor blockers against ventilator-induced diaphragm dysfunction in rats. Clin Transl Sci 2021;14(02): 481–486
- 15 Das K, Kumar S, Bindra A, Kumar N, Pathak S. Ultrasound assessment of diaphragm function in traumatic brain injury: a prospective observational study. J Neuroanaesthesiol Crit Care 2023;10(03): 183–188
- 16 Panelli A, Verfuß MA, Dres M, Brochard L, Schaller SJ. Phrenic nerve stimulation to prevent diaphragmatic dysfunction and ventilator-induced lung injury. Intensive Care Med Exp 2023;11 (01):94–106