



# Effects of Sleep-Disordered Breathing on Daytime Brain Activity in Community-Dwelling Older Adults

Hiroki Shibata<sup>1</sup> Akiko Noda<sup>1</sup> Yuanjie Mao<sup>2</sup> Kunihiro Iwamoto<sup>3</sup> Masato Okuda<sup>1</sup> Ippei Okada<sup>3</sup>  
Seiko Miyata<sup>3</sup> Toshiaki Taoka<sup>4</sup> Fumihiko Yasuma<sup>5</sup>

<sup>1</sup> Department of Biomedical Sciences, Graduate School of Life and Health Sciences, Chubu University, Kasugai-shi, Aichi, Japan

<sup>2</sup> Heritage College of Osteopathic Medicine, Ohio University, Athens, Ohio, United States

<sup>3</sup> Department of Psychiatry, Nagoya University Graduate School of Medicine, NAGOYA, Aichi, Japan

<sup>4</sup> Department of Innovative Biomedical Visualization, Graduate School of Medicine, Nagoya University, Nagoya, Aichi, Japan

<sup>5</sup> Department of Biomedical Sciences, College of Life and Health Sciences, Chubu University, Kasugai-shi, Aichi, Japan

**Address for correspondence** Akiko Noda  
(e-mail: anoda@isc.chubu.ac.jp).

Sleep Sci

## Abstract

**Introduction** Sleep-disordered breathing (SDB) is associated with an increased risk of cardiovascular diseases. The present study aimed to examine the influence of SDB on daytime brain activity in the community-dwelling older adults.

**Material and Methods** There were 81 consecutive volunteers aged 60 years or older (mean age  $70.5 \pm 4.8$  years) who participated in the present study. Daytime brain activity was assessed by measuring the peak cortical oxygenated hemoglobin (OxyHb) levels and area under the near-infrared spectroscopy (NIRS) curve. The respiratory event index (REI) and 3% oxygen desaturation index (3%ODI) were evaluated using a home sleep-apnea test.

**Results** The peak OxyHb and area under the NIRS curve were significantly lower in the participants with  $REI \geq 15/h$  than those with  $REI < 15/h$ . The body mass index (BMI), REI, 3%ODI, and Epworth sleepiness scale (ESS) scores were significantly correlated with peak OxyHb levels (BMI:  $r = -0.202$ ,  $p = 0.035$ ; REI:  $r = -0.307$ ,  $p = 0.003$ ; 3%ODI:  $r = -0.321$ ,  $p = 0.003$ ; and ESS score:  $r = -0.287$ ,  $p = 0.005$ ). Also, the BMI, REI, and 3%ODI were significantly correlated with the area under the NIRS curve (BMI:  $r = -0.306$ ,  $p = 0.003$ ; REI:  $r = -0.326$ ,  $p = 0.002$ ; and 3%ODI:  $r = -0.313$ ,  $p = 0.002$ ), and BMI was a significant factor associated with the area under the NIRS curve.

**Conclusions** Brain activity during wakefulness was associated with SDB and BMI severity. A simple NIRS may yield unique information for characterizing the decline in daytime brain activity of the community-dwelling older adults.

## Keywords

- ▶ brain activity
- ▶ sleep-disordered breathing
- ▶ daytime sleepiness
- ▶ older adults
- ▶ near-infrared spectroscopy

received  
April 1, 2023  
accepted  
February 1, 2024

DOI <https://doi.org/10.1055/s-0044-1782627>.  
ISSN 1984-0659.

© 2024, Brazilian Sleep Association. All rights reserved.  
This is an open access article published by Thieme under the terms of the Creative Commons Attribution-NonDerivative-NonCommercial-License, permitting copying and reproduction so long as the original work is given appropriate credit. Contents may not be used for commercial purposes, or adapted, remixed, transformed or built upon. (<https://creativecommons.org/licenses/by-nc-nd/4.0/>)  
Thieme Revinter Publicações Ltda., Rua do Matoso 170, Rio de Janeiro, RJ, CEP 20270-135, Brazil

## Introduction

The prevalence of sleep-disordered breathing (SDB) was estimated as high as 20 to 40% in the geriatric population, and it was expected to increase with the population's aging.<sup>1,2</sup> This disorder leads to a potential risk of cerebrovascular and cardiovascular diseases,<sup>3</sup> and might lead to hypoxemia, pressure swings of the thorax, frequent arousals, sleep fragmentation, and activation of the sympathetic nervous system.<sup>4-6</sup> Chronic intermittent episodes of hypoxia could induce oxidative stress, inflammation of the whole body, and, possibly, degeneration of the central nervous system.<sup>7</sup> Moreover, quality of life was impaired by the coexisting SDB, which decreased mental/physical activity.<sup>8</sup> Therefore, an early detection and appropriate treatment of SDB have beneficial effects on the cognitive function of older adults.

According to the previous studies using the functional magnetic resonance imaging (fMRI), the broad functional connectivity was reduced among the frontal, cingulate, temporal, parietal, occipital, and cerebellar regions of the brain in patients with SDB.<sup>9</sup> Near-infrared spectroscopy (NIRS) with high temporal resolution is a noninvasive method for assessing the brain activity, which measured the relative changes in oxygenated hemoglobin (OxyHb) of the cerebral cortex.<sup>10</sup> In a previous study, we already reported that short sleep duration had a negative impact on the cortical oxygenation during a word fluency task evaluated using the NIRS.<sup>11,12</sup> Since the target population of the present study is community-dwelling older adults, we have utilized the NIRS to measure brain activity, as it is simpler, less expensive, and less invasive than fMRI.

The present study aims to clarify the relationship between brain activity and SDB in the target population.

## Material and Methods

### Participants

A total of 81 consecutive volunteers aged  $\geq 60$  years (51 men, 30 women; mean age  $70.5 \pm 4.8$  years) were enrolled in the present study. The participants had no history of myocardial infarction, angina pectoris, heart failure, cerebral infarction, cerebral hemorrhage, chronic obstructive pulmonary disease, psychiatric disorders, or the use of antidepressants, benzodiazepines, or current medications for sleep problems. We used a questionnaire to collect the following data: age; body mass index (BMI); smoking status; alcohol intake; history of hypertension, diabetes mellitus, and/or hyperlipidemia; and current use of medications. Active smokers were defined as the participants who were currently smoking or those who had quit smoking for  $< 4$  years prior.<sup>13</sup> Alcohol intake was defined as an individual who drank regularly.<sup>14</sup> Systolic (SBP) and diastolic blood pressure (DBP) were measured using a plethysmograph (BP-203RPEIII, Omron Colin Co., Ltd., Tokyo, Japan) prior to the NIRS measurements. The participants with SBP  $\geq 140$  mm Hg or DBP  $\geq 90$  mm Hg, or those undergoing antihypertensive therapy, were considered to have hypertension.<sup>15</sup> Diabetes mellitus and

hyperlipidemia were defined by the use of oral hypoglycemic and lipid-lowering agents, respectively.

The protocol for the present study was approved by the Ethics Committee of the Chubu University (number 270098). After explaining the nature of the study and the procedures involved, the written informed consents were obtained from all participants.

### Questionnaires

#### Epworth Sleepiness Scale (ESS)

Subjective excessive daytime sleepiness was evaluated using the Epworth sleepiness scale (ESS).<sup>16</sup> In this questionnaire, the participants used a four-point scale to rate their chances of dozing off in eight different situations, all of which can be encountered in daily life. The total ESS score was the sum of all responses, ranging from 0 to 24 points.

#### Pittsburgh Sleep Quality Index

Subjective sleep quality over the prior month was assessed using the Pittsburgh sleep quality index (PSQI),<sup>17</sup> which contained 19 items in 7 component domains: subjective sleep quality, sleep latency, sleep duration, habitual sleep efficiency, sleep disturbances, use of sleeping medication, and daytime dysfunction. The 19 self-rated items were combined to form 7 "component" scores, each of which had a range from 0 to 3 points. In all cases, a score of 0 indicated no difficulty, while a score of 3 indicated severe difficulty. The 7 component scores were then totaled to yield a global PSQI score, ranging from 0 to 21 points.

### Near-Infrared Spectroscopy (NIRS) During Word Frequency Task


Relative concentrations of OxyHb were measured using the 2-channel NIRS recorder (HOT121B; Hitachi High-Technologies Co., Tokyo, Japan), while the participant underwent a word fluency task. The NIRS probes (3 cm distance between the emitter and detector probes) were placed on the left and right frontal areas, including Fp1 and Fp2, according to the international 10 to 20 electroencephalography system of the electroencephalography. The absorption of near-infrared light was measured with a temporal resolution of 0.1 second.

The word frequency task was performed as previously described.<sup>18</sup> In brief, the 30-second pretask involved repeating the vowels 'a, e, i, o, u', and then speaking out any random words starting with the three initial Japanese syllables in 20 seconds each. This was followed by a 60-second post-task, which again involved repeating the vowels, and a 70-second posttask (**- Fig. 1**). The values recorded from both channels in the frontal area were then averaged to obtain the results.

Peak OxyHb was measured during the word frequency task. The area under the NIRS curve was also measured using the trapezium rule, which approximated this area, as described by the function  $f(x)$ .

### Home Sleep Apnea Test (HSAT)

The participants were screened for SDB using a portable device (SAS-2100, Nihon Kohden Corp., Tokyo, Japan), in



**Fig. 1** The NIRS curve of a representative case. During the word fluency task, the area under the NIRS curve was measured using the trapezium rule, which approximated that area to measure the integral. Peak OxyHb was also measured.

**Table 1** Baseline characteristics and sleep/NIRS parameters by gender.

	Total (n = 81)	Men (n = 51)	Women (n = 30)	p-value
<b>Demographics</b>				
Age (years)	70.5 ± 4.8	70.1 ± 4.1	71.2 ± 5.9	0.377
Height (cm)	161.0 ± 8.6	165.6 ± 6.7	153.2 ± 5.1	< 0.001
Weight (kg)	60.0 ± 10.4	64.2 ± 9.5	53.0 ± 8.1	< 0.001
BMI (kg/m <sup>2</sup> )	23.0 ± 3.1	23.2 ± 3.2	22.8 ± 2.9	0.591
SBP (mmHg)	141.5 ± 15.4	142.6 ± 13.9	139.6 ± 17.6	0.399
DBP (mmHg)	83.6 ± 9.1	84.9 ± 8.4	81.4 ± 9.9	0.099
<b>Risk factors</b>				
Smoking (%)	33.3	49.0	6.7	< 0.001
Alcohol intake (%)	59.2	74.5	33.3	< 0.001
Hypertension (%)	54.3	54.9	53.3	0.891
Diabetes mellitus (%)	4.9	43.9	6.7	0.474
Hyperlipidemia (%)	16.0	7.8	30.0	0.011
<b>Sleep parameters</b>				
Questionnaires				
Sleep duration (min)	428.3 ± 73.4	434.2 ± 79.8	418.4 ± 60.9	0.354
ESS score	5.6 ± 3.9	5.5 ± 3.9	5.9 ± 3.9	0.661
PSQI score	5.5 ± 2.9	5.5 ± 3.1	5.5 ± 2.6	0.988
Home sleep apnea test				
REI (/h)	9.7 ± 9.5	10.3 ± 10.1	8.6 ± 8.2	0.425
3%ODI (/h)	14.1 ± 11.1	14.9 ± 11.7	12.8 ± 10.1	0.402
Minimum SpO <sub>2</sub> (%)	86.2 ± 5.2	86.3 ± 5.2	86.2 ± 5.3	0.986
<b>NIRS parameters</b>				
Peak OxyHb (mM × mm)	0.316 ± 0.260	0.302 ± 0.255	0.339 ± 0.270	0.549
Area under the NIRS curve	102.6 ± 144.5	78.87 ± 135.3	143.0 ± 152.9	0.053

**Abbreviations:** BMI, body mass index; DBP, diastolic blood pressure; ESS, Epworth sleepiness scale; NIRS, near-infrared spectroscopy; ODI, oxygen desaturation index; OxyHb, oxygenated hemoglobin; PSQI, Pittsburgh sleep quality index; REI, respiratory event index; SBP, systolic blood pressure. **Notes:** Data are expressed as mean ± SD.

which a nasal pressure sensor and a pulse oximeter were used to record airflow, pulse wave, and oxygen saturation (SpO<sub>2</sub>), respectively. They were instructed on how to wear and use the device at home. We calculated the respiratory event index (REI) as the total number of apnea and hypopnea events occurring per hour for the total duration of the recording time. The number of events per hour with oxygen desaturation  $\geq 3\%$  (3% oxygen desaturation index: 3%ODI). The minimum SpO<sub>2</sub> values were also evaluated.

### Statistical Analyses

All data are presented as mean  $\pm$  standard deviation (SD). We compared data on the smoking status, alcohol intake, hypertension, diabetes mellitus, hyperlipidemia, medication use, ESS score, PSQI score, HSAT results, and NIRS parameters between the groups (men vs. women, REI  $\geq 15$ /h vs.  $< 15$ /h,<sup>19,20</sup> and 3%ODI  $\geq 15$ /h vs.  $< 15$ /h) using the chi-squared test or non-paired *t*-test. Based on the results of the preliminary test, we calculated the number of subjects

required to verify this study and the power. The Pearson correlation analysis was performed to evaluate the relationships between the sleep parameters and NIRS.

Moreover, a multiple linear regression analysis was performed to examine the association of NIRS parameters (peak OxyHb and area under the NIRS curve) with age, BMI, sex, smoking status, sleep duration, REI, 3%ODI, ESS score, and PSQI score. Significant factors were obtained from the correlation analysis. Probability values lower than 0.05 were considered statistically significant. All statistical analyses were performed using the Statistical Package Social Sciences (SPSS, IBM Corporation, Armonk, NY, USA) version 25.0.

### Results

The baseline characteristics of the study participants are shown in ►Table 1. Men had a significantly higher prevalence of smoking and alcohol use, and a significantly lower incidence of hyperlipidemia than women. There were,

**Table 2** Baseline characteristics and sleep/NIRS parameters by respiratory event index.

	REI < 15/h (n = 61)	REI $\geq$ 15/h (n = 20)	p-value
<b>Demographics</b>			
Age (years)	70.8 $\pm$ 4.8	69.8 $\pm$ 4.8	0.445
Men (%)	60.7	70.0	0.453
Height (cm)	160.5 $\pm$ 8.6	162.4 $\pm$ 8.5	0.396
Weight (kg)	57.3 $\pm$ 8.5	68.4 $\pm$ 11.6	< 0.001
BMI (kg/m <sup>2</sup> )	22.2 $\pm$ 2.5	25.4 $\pm$ 3.7	0.002
SBP (mmHg)	140.1 $\pm$ 16.0	145.7 $\pm$ 12.6	0.158
DBP (mmHg)	82.7 $\pm$ 9.5	86.2 $\pm$ 7.2	0.134
<b>Risk factors</b>			
Smoking (%)	29.5	45.0	0.202
Alcohol intake (%)	62.3	50.0	0.331
Hypertension (%)	54.1	55.0	0.944
Diabetes mellitus (%)	6.6	0.0	0.314
Hyperlipidemia (%)	18.0	10.0	0.322
<b>Sleep parameters</b>			
<b>Questionnaires</b>			
Sleep duration (min)	426.5 $\pm$ 76.3	433.9 $\pm$ 65.0	0.697
ESS score	5.0 $\pm$ 3.5	7.5 $\pm$ 4.5	0.014
PSQI score	5.5 $\pm$ 2.8	5.5 $\pm$ 3.3	0.991
Home sleep apnea test			
3%ODI (/h)	8.9 $\pm$ 5.4	30.1 $\pm$ 8.5	< 0.001
Minimum SpO <sub>2</sub> (%)	87.6 $\pm$ 4.6	82.1 $\pm$ 4.6	< 0.001
<b>NIRS parameters</b>			
Peak OxyHb (mM $\times$ mm)	0.357 $\pm$ 0.280	0.191 $\pm$ 0.124	< 0.001
Area under the NIRS curve	123.1 $\pm$ 149.0	40.29 $\pm$ 111.6	0.025

**Abbreviations:** BMI, body mass index; DBP, diastolic blood pressure; ESS, Epworth sleepiness scale; NIRS, near-infrared spectroscopy; ODI, oxygen desaturation index; OxyHb, oxygenated hemoglobin; PSQI, Pittsburgh sleep quality index; REI, respiratory event index; SBP, systolic blood pressure.  
**Notes:** Data are expressed as mean  $\pm$  SD.

however, no significant differences in the sleep and NIRS parameters between men and women.

Body weight, BMI, and ESS score were significantly higher in the participants with  $REI \geq 15/h$  than those  $< 15/h$ . Peak OxyHb and area under the NIRS curve significantly decreased in the participants with  $REI \geq 15/h$  than those  $< 15/h$  (► **Table 2**). The number of subjects required to verify this study was 32 cases in one group and 64 cases in the two groups with the power of 77.6%.

Body weight, BMI, and ESS score significantly increased in participants with  $3\%ODI \geq 15/h$  than those  $< 15/h$ . Peak OxyHb, and the area under the NIRS curve significantly decreased in the participants with  $3\%ODI \geq 15/h$  than those with  $< 15/h$  (► **Table 3**). The number of subjects required to verify this study was 32 cases in one group and 64 cases in the two groups with the power of 86.7%.

The BMI ( $r = -0.202$ ,  $p = 0.035$ ), REI ( $r = -0.307$ ,  $p = 0.003$ ),  $3\%ODI$  ( $r = -0.321$ ,  $p = 0.002$ ), and ESS score ( $r = -0.287$ ,  $p = 0.005$ ) were significantly correlated with peak OxyHb

levels. Also, the BMI ( $r = -0.306$ ,  $p = 0.002$ ), REI ( $r = -0.326$ ,  $p = 0.002$ ), and  $3\%ODI$  ( $r = -0.322$ ,  $p = 0.002$ ) were significantly correlated with the area under the NIRS curve. Multiple regression analysis revealed that BMI was a significant factor for the area under the NIRS curve ( $\beta = -0.247$ ,  $p = 0.044$ ) (► **Table 4**).

## Discussion

We found that the peak OxyHb and area under the NIRS curve were decreased with increasing REI and  $3\%ODI$ , and that increases in the ESS score and BMI were associated with the REI and  $3\%ODI$ . These results suggest that SDB may play a significant role in the decrease of daytime brain activity of community-dwelling older adults.

A previous fMRI study demonstrated that brain activity decreased significantly in the 24 patients with SDB than in the 24 healthy volunteers,<sup>21</sup> which seemed consistent with the findings of our present study. From a methodological

**Table 3** Baseline characteristics and sleep/NIRS parameters by 3% oxygen desaturation index.

	3%ODI < 15/h (n = 52)	3%ODI ≥ 15/h (n = 29)	p-value
<b>Demographics</b>			
Age (years)	70.9 ± 4.9	69.8 ± 4.7	0.313
Men (%)	59.6	69.0	0.403
Height (cm)	160.6 ± 7.9	161.6 ± 9.7	0.620
Weight (kg)	57.0 ± 8.4	65.6 ± 11.6	< 0.001
BMI (kg/m <sup>2</sup> )	22.1 ± 2.5	24.7 ± 3.7	< 0.001
SBP (mmHg)	139.7 ± 17.0	144.7 ± 11.5	0.161
DBP (mmHg)	82.3 ± 9.9	85.9 ± 7.1	0.088
<b>Risk factors</b>			
Smoking (%)	28.8	41.4	0.251
Alcohol intake (%)	59.6	58.6	0.930
Hypertension (%)	44.8	54.3	0.200
Diabetes mellitus (%)	5.8	3.4	0.548
Hyperlipidemia (%)	21.2	6.9	0.083
<b>Sleep parameters</b>			
<b>Questionnaires</b>			
Sleep duration (min)	427.0 ± 72.3	430.8 ± 76.4	0.822
ESS score	4.7 ± 3.6	7.2 ± 4.0	0.005
PSQI score	5.3 ± 2.6	5.9 ± 3.4	0.312
<b>Home sleep apnea test</b>			
REI (/h)	4.1 ± 2.9	19.6 ± 8.9	< 0.001
Minimum SpO <sub>2</sub> (%)	88.3 ± 4.4	82.6 ± 4.5	< 0.001
<b>NIRS parameters</b>			
Peak OxyHb (mM × mm)	0.378 ± 0.295	0.204 ± 0.119	< 0.001
Area under the NIRS curve	135.1 ± 154.8	44.39 ± 102.9	0.006

**Abbreviations:** BMI, body mass index; DBP, diastolic blood pressure; ESS, Epworth Sleepiness Scale; NIRS, near-infrared spectroscopy; ODI, oxygen desaturation index; OxyHb, oxygenated hemoglobin; PSQI, Pittsburgh Sleep Quality Index; REI, respiratory event index; SBP, systolic blood pressure. **Notes:** Data are expressed as mean ± SD.

**Table 4** Relationship between NIRS parameters and baseline/sleep parameters.

	Simple correlation analysis		Multiple regression analysis		Simple correlation analysis		Multiple regression analysis	
	<i>r</i>	<i>p</i> -value	$\beta$	<i>p</i> -value	<i>r</i>	<i>p</i> -value	$\beta$	<i>p</i> -value
	Peak OxyHb				Area under the NIRS curve			
Age	-0.121	0.141	-0.182	0.115	-0.064	0.286	-0.155	0.168
BMI	-0.202	0.035	-0.104	0.402	-0.306	0.003	-0.247	0.044
Sex			-0.023	0.850		0.027	-0.121	0.306
Smoking			-0.115	0.361		0.008	-0.181	0.143
Sleep duration	-0.011	0.460	0.003	0.980	-0.124	0.135	-0.061	0.585
REI	-0.307	0.003	0.265	0.559	-0.326	0.002	-0.079	0.858
3%ODI	-0.321	0.002	-0.458	0.313	-0.322	0.002	-0.079	0.859
ESS score	-0.287	0.005	-0.216	0.066	-0.148	0.094	-0.064	0.574
PSQI score	-0.081	0.237	-0.006	0.956	-0.016	0.444	0.033	0.770

**Abbreviations:** BMI, body mass index; ESS, Epworth sleepiness scale; NIRS, near-infrared spectroscopy; ODI, oxygen desaturation index; OxyHb, oxygenated hemoglobin; PSQI, Pittsburgh sleep quality index; REI, respiratory event index.

viewpoint, the NIRS is a simple technology for assessing daytime brain activity through OxyHb of the cerebral cortex. In contrast, the fMRI was time-consuming and expensive,<sup>10</sup> making it difficult to assess the brain's neural function in community-dwelling older adults. Our previous NIRS study demonstrated that the cortical OxyHb response was blunted during word fluency tasks, which was associated with the short sleep duration in elderly volunteers.<sup>22</sup> This test, with its methodological and financial advantages over fMRI, could provide a novel insight into the brain activity of community-dwelling older adults.

Increases in REI and 3%ODI were found to be associated with an increase in the ESS score. Intermittent episodes of hypoxia and frequent arousals during sleep in patients with SDB could cause the symptoms of excessive daytime sleepiness, presumably.<sup>23</sup> For example, we previously showed that the arousal index was the most reliable predictor for the occurrence of automobile accidents, and that their incidence, that of near-misses, as well as scores of daytime sleepiness scores, were significantly correlated with the severity of nocturnal hypoxemia.<sup>24</sup> Moreover, we revealed that sleep fragmentation due to SDB led to an augmented sympathetic nerve activity during nighttime sleep.<sup>5,6</sup> Therefore, the symptoms of excessive daytime sleepiness can be attributed to the impaired brain activity during daytime wakefulness.

Sleep disturbances are common in aging and there is an increase in related disorders in pathological aging. These disturbances are characterized by decreased sleep duration, quality, and efficiency, and increased sleep fragmentation.<sup>25,26</sup> However, in our study, the peak OxyHb and area under the NIRS curve were not correlated with the PSQI score. In a previous study, we showed that a long sleep time and an irregular sleep-wake rhythm could have adverse effects on executive function and working memory in older people.<sup>27</sup> Hence, it might be difficult to evaluate the sleep problems associated with brain activity in older adults using PSQI alone. Mecca et al. demonstrated that there was no association between sleep disturbance and im-

paired cognitive functioning.<sup>28</sup> Kyle et al. have reported higher cognitive functioning in a large sample of participants with insomnia.<sup>29</sup> These previous reports seem support our findings.

In the present study, we showed that an increase in BMI was associated with an increase in REI and 3%ODI, and that BMI was a significant independent factor for the area under the NIRS curve. These results suggest that obesity might affect the occurrences of SDB and oxygen desaturation in the cerebral cortex, the latter of which was noted in the word frequency task during daytime wakefulness. A cohort study of men and women with a mean-age of 62 years reported a higher probability of association between gaining body weight and development of SDB.<sup>30</sup> Regarding this role, a previous study reported that apnea and hypopnea indices had worse results on patients with enhanced esophageal pressure, oxygen desaturation, and severity of obesity.<sup>31</sup> Studies have also shown that obesity can cause neuroinflammation, insulin resistance, and mitochondrial perturbations in the brain, which might impair cognitive function.<sup>32</sup> Thus, SDB aggravates oxygen desaturation during nighttime sleep in moderate to severe obesity, which in turn could jeopardize the reduced brain activity during daytime wakefulness.

There are several limitations of this study. First, neuroimaging techniques such as the fMRI and positron emission tomography can detect morphological changes in the cerebral vasculature,<sup>33</sup> whereas the NIRS is limited to OxyHb changes on the cortical surface. Second, the number of participants was relatively small, but the number of necessary subjects was satisfied at 81 subjects, considering the power. To clarify the effects of SDB on the daytime brain activity, an intervention study encompassing a large population is necessary in the future.

## Conclusions

We found that SDB with symptomatic daytime sleepiness adversely affected brain activity. The simple NIRS should



prove valuable for determining the decline in daytime brain activity and for detecting SDB in community-dwelling older adults.

#### Funding Source

The study was supported by the Japan Society for the Promotion of Science (KAKENHI Grant Number 25282210, 21K07563).

#### Conflict of Interests

The authors have no conflict of interests to declare.

#### References

- Osorio RS, Martínez-García MÁ, Rapoport DM. Sleep apnoea in the elderly: a great challenge for the future. *Eur Respir J* 2021;59(04):2101649
- Senaratna CV, Perret JL, Lodge CJ, et al. Prevalence of obstructive sleep apnea in the general population: A systematic review. *Sleep Med Rev* 2017;34:70–81
- Gottlieb DJ, Punjabi NM. Diagnosis and management of obstructive sleep apnea: a review. *JAMA* 2020;323(14):1389–1400
- Somers VK, Dyken ME, Clary MP, Abboud FM. Sympathetic neural mechanisms in obstructive sleep apnea. *J Clin Invest* 1995;96(04):1897–1904
- Noda A, Yasuma F, Okada T, Yokota M. Circadian rhythm of autonomic activity in patients with obstructive sleep apnea syndrome. *Clin Cardiol* 1998;21(04):271–276
- Noda A, Hayano J, Ito N, et al. Very low frequency component of heart rate variability as a marker for therapeutic efficacy in patients with obstructive sleep apnea: Preliminary study. *J Res Med Sci* 2019;24:84
- Snyder B, Shell B, Cunningham JT, Cunningham RL. Chronic intermittent hypoxia induces oxidative stress and inflammation in brain regions associated with early-stage neurodegeneration. *Physiol Rep* 2017;5(09):e13258
- Baldwin CM, Griffith KA, Nieto FJ, et al. The association of sleep-disordered breathing and sleep symptoms with quality of life in the Sleep Heart Health Study. *Sleep* 2001;24(01):96–105
- Park HR, Cha J, Joo EY, Kim H. Altered cerebrocerebellar functional connectivity in patients with obstructive sleep apnea and its association with cognitive function. *Sleep* 2022;45(01):zsab209
- Cui X, Bray S, Bryant DM, et al. A quantitative comparison of NIRS and fMRI across multiple cognitive tasks. *Neuroimage* 2011;54(04):2808–2821
- Miyata S, Noda A, Iwamoto K, et al. Impaired cortical oxygenation is related to mood disturbance resulting from three nights of sleep restriction. *Sleep Biol Rhythms* 2015;13:387–394
- Kato K, Iwamoto K, Kawano N, et al. Differential effects of physical activity and sleep duration on cognitive function in young adults. *J Sport Health Sci* 2018;7(02):227–236
- Kondo T, Osugi S, Shimokata K, et al. Smoking and smoking cessation in relation to all-cause mortality and cardiovascular events in 25,464 healthy male Japanese workers. *Circ J* 2011;75(12):2885–2892
- Cho Y, Shin SY, Won S, et al. Alcohol intake and cardiovascular risk factors: A Mendelian randomisation study. *Sci Rep* 2015;5:18422
- Umemura S, Arima H, Arima S, et al. The Japanese Society of Hypertension Guidelines for the Management of Hypertension (JSH 2019). *Hypertens Res* 2019;42(09):1235–1481
- Takegami M, Suzukamo Y, Wakita T, et al. Development of a Japanese version of the Epworth Sleepiness Scale (JESS) based on item response theory. *Sleep Med* 2009;10(05):556–565
- Doi Y, Minowa M, Uchiyama M, et al. Psychometric assessment of subjective sleep quality using the Japanese version of the Pittsburgh Sleep Quality Index (PSQI-J) in psychiatric disordered and control subjects. *Psychiatry Res* 2000;97(2-3):165–172
- Nishimura Y, Tani H, Fukuda M, et al. Frontal dysfunction during a cognitive task in drug-naive patients with panic disorder as investigated by multi-channel near-infrared spectroscopy imaging. *Neurosci Res* 2007;59(01):107–112
- Caples SM, Anderson WM, Calero K, et al. Use of polysomnography and home sleep apnea tests for the longitudinal management of obstructive sleep apnea in adults: an American Academy of Sleep Medicine clinical guidance statement. *J Clin Sleep Med* 2021;17(06):1287–1293
- Kapur VK, Auckley DH, Chowdhuri S, et al. Clinical practice guideline for diagnostic testing for adult obstructive sleep apnea: an American Academy of Sleep Medicine clinical practice guideline. *J Clin Sleep Med* 2017;13(03):479–504
- Peng DC, Dai XJ, Gong HH, et al. Altered intrinsic regional brain activity in male patients with severe obstructive sleep apnea: a resting-state functional magnetic resonance imaging study. *Neuropsychiatr Dis Treat* 2014;10:1819–1826
- Kato K, Miyata S, Ando M, et al. Influence of sleep duration on cortical oxygenation in elderly individuals. *Psychiatry Clin Neurosci* 2017;71(01):44–51
- Lal C, Weaver TE, Bae CJ, Strohl KP. Mechanisms and Clinical Management. Excessive daytime sleepiness in obstructive sleep apnea. mechanisms and clinical management. *Ann Am Thorac Soc* 2021;18(05):757–768
- Noda A, Yasuma F, Miyata S, et al. Sleep fragmentation and risk of automobile accidents in patients with obstructive sleep apnea. *Health* 2019;11(02):171–181
- Gadie A, Shafto M, Leng Y, Kievit RAC. How are age-related differences in sleep quality associated with health outcomes? An epidemiological investigation in a UK cohort of 2406 adults. *BMJ Open* 2017;7(07):e014920
- Mander BA, Winer JR, Walker MP. Sleep and human aging. *Neuron* 2017;94(01):19–36
- Okuda M, Noda A, Iwamoto K, et al. Effects of long sleep time and irregular sleep-wake rhythm on cognitive function in older people. *Sci Rep* 2021;11(01):7039
- Mecca AP, Michalak HR, McDonald JW, et al. The Alzheimer's Disease Neuroimaging Initiative (ADNI) Sleep disturbance and the risk of cognitive decline or clinical conversion in the ADNI cohort. *Dement Geriatr Cogn Disord* 2018;45(3-4):232–242
- Kyle SD, Sexton CE, Feige B, et al. Sleep and cognitive performance: cross-sectional associations in the UK Biobank. *Sleep Med* 2017;38:85–91
- Newman AB, Foster G, Givelber R, et al. Progression and regression of sleep-disordered breathing with changes in weight: the Sleep Heart Health Study. *Arch Intern Med* 2005;165(20):2408–2413
- Itasaka Y, Miyazaki S, Ishikawa K, Togawa K. The influence of sleep position and obesity on sleep apnea. *Psychiatry Clin Neurosci* 2000;54(03):340–341
- Lee TH, Yau SY. From obesity to hippocampal neurodegeneration: pathogenesis and non-pharmacological interventions. *Int J Mol Sci* 2020;22(01):201
- Detre JA, Wang J. Technical aspects and utility of fMRI using BOLD and ASL. *Clin Neurophysiol* 2002;113(05):621–634