SYNLETT Spotlight 61

This feature focuses on a reagent chosen by a postgraduate, highlighting the uses and preparation of the reagent in current research

PCC: Novel Oxidation Reactions

Compiled by Rodney A. Fernandes

Rodney A. Fernandes was born in Goa (India) in 1972. He received his MSc in Organic Chemistry (1997) from Goa University and PhD in Chemistry (2003) from Pune University (India) under the tutelage of Dr. Pradeep Kumar. His main interest is in the asymmetric synthesis of bioactive natural products and organic transformations.

SRF, Organic Chemistry: Technology, National Chemical Laboratory, Pune 411 008, India E-mail: fernandesrod@yahoo.co.in

Introduction

Until the studies of Corey and co-workers (1975),¹ the reactivity of pyridinium chlorochromate (PCC) had been investigated little. The first report on the chlorochromate anion dates back to 1833.² In 1899, the first synthesis of PCC was achieved.³ PCC is prepared easily and safely.¹ It shows a high capability to convert primary alcohols exclusively to aldehydes. Mere filtration, washing of the residue with diethyl ether, and concentration of the filtrate furnish virtually pure product. PCC shows a slightly acidic character and for acid-labile groups the reaction could be buffered with NaOAc. The applications of PCC are not limited to oxidation reactions, but several rearrangements and useful conversions are mediated by PCC, which makes it a versatile reagent in organic synthesis.



Preparation

The addition of CrO_3 to 6 N HCl furnishes the unstable chlorochromic acid and subsequent addition of pyridine at 0 °C immediately gives a yellow-orange solid of PCC which is not appreciably hygroscopic.¹

Abstracts



Synlett 2003, No. 5, Print: 10 04 2003. Art Id.1437-2096,E;2003,0,05,0741,0742,ftx,en;V06502ST.pdf. © Georg Thieme Verlag Stuttgart · New York ISSN 0936-5214

R-CH₂CHO

RCHO

сно

RCHO

PCC

PCC

PCC

(D) Hydroboration of 1-alkenes give trialkyl boranes which on subsequent oxidation with PCC furnish aldehydes.^{7a} PCC oxidation of borate esters or boroxines also give aldehydes.^{7b,c}

(E) PCC oxidation of tertiary 2-alkyl cyclopropyl carbinols furnished transposed β , γ -enones, making the overall process a synthetically useful method for 1,4-carbonyl transposition.⁸

(F) PCC can release quinone from hydroquinone silyl ethers.9



(R-CH2-CH2)3-B

(RCH₂O)₃B

1/3(RCH₂OBO)₃

 R^1

H₂B·SMe₂

H₃B·SMe₂

H₃B·SMe₂

R-CH=CH₂

R-CH₂OH

R-CO₂H

(G) The allylic alcohols obtained from lithiated 1,4-dioxene and ketones or aldehydes undergo regiospecific oxidative cleavage with PCC at the dioxene double bond to furnish, after saponification, α -hydroxy acids. Extension of this reaction to α , β -unsaturated ketones affords α -keto acids.¹⁰



(H) Specific oxidative cleavage of allylic and benzylic ethers is known with PCC. 11



PCC

OH

R = alkyl, aryl

(I) Oxidation of homoallylic alcohol with excess of PCC gave products arising from alcohol oxidation, double bond migration with isomerization from the *cis*- to the *trans*-olefin and subsequent allylic oxidation. Thus, this constitutes a synthetically useful conversion of homoallylic alcohols to 1,4-dicarbonyl-2-ene compounds.¹²

References

- (1) Corey, E. J.; Suggs, J. W. Tetrahedron Lett. 1975, 2647.
- (2) Sisler, H. H. Inorg. Synth. 1946, 2, 208.
- (3) Meyer, R. J.; Best, Z. Anorg. Allgem. Chem. 1899, 22, 192.
- (4) (a) Babler, J. H.; Coghlan, M. J. Synth. Commun. 1976, 6, 469. (b) Sundararaman, P.; Herz, W. J. Org. Chem. 1977, 42, 806. (c) Sundararaman, P.; Herz, W. J. Org. Chem. 1977, 42, 813. (d) Dauben, W. G.; Michno, D. M. J. Org. Chem. 1977, 42, 682.
- (5) Piancatelli, G.; Scettri, A.; D'Auria, M. *Tetrahedron Lett.* 1977, 2199.
- (6) (a) Maloney, J. R.; Lyle, R. E.; Saavedra, J. E.; Lyle, G. G. Synthesis 1978, 212. (b) Drabowicz, J. Synthesis 1980, 125.
- (7) (a) Rao, C. G.; Kulkarni, S. U.; Brown, H. C. J. Organomet. Chem. 1979, 172, C20. (b) Brown, H. C.; Kulkarni, S. U.; Rao, C. G. Synthesis 1979, 702. (c) Brown, H. C.; Rao, C. G.; Kulkarni, S. U. Synthesis 1979, 704.

СНО

- (8) Nakai, T.; Wada, E.; Okawara, M. J. Org. Chem. 1979, 44, 2952.
- (9) Willis, J. P.; Gogins, K. A. Z.; Miller, L. L. J. Org. Chem. 1981, 46, 3215.
- (10) Fetizon, M.; Goulaouic, P.; Hanna, I. *Tetrahedron Lett.* 1988, 29, 6261.
- (11) Cossy, J.; Bouzbouz, S.; Lachgar, M.; Hakiki, A.; Tabyaouic, B. *Tetrahedron Lett.* **1998**, *39*, 2561.
- (12) Fernandes, R. A.; Kumar, P. *Tetrahedron Lett.* 2003, 44, 1275.

Synlett 2003, No. 5, 741–742 $\,$ ISSN 0936-5214 $\,$ © Thieme Stuttgart \cdot New York