


**K. Ragnath, G. K. Anagnostopoulos**  
Wolfson Digestive Diseases Centre,  
Queen's Medical Centre, University of  
Nottingham, Nottingham,  
United Kingdom

**Competing interests:** Not declared


### Corresponding Author

**G. K. Anagnostopoulos, M. D.**

Wolfson Digestive Diseases Centre,  
Queen's Medical Centre,  
University of Nottingham  
29A Newcastle Drive  
The Park  
Nottingham NG7 1AA  
United Kingdom  
Fax: +44-7834-776320  
E-mail: gkanagnostopoulos@yahoo.gr



**Figure 1** Conventional endoscopic view of the prepyloric region of the stomach, showing a well-circumscribed area of reddened mucosa which was interpreted as "antral gastritis".



**Figure 2** The same region visualised using a high-resolution zoom endoscope (Olympus GIF-Q240Z; Olympus Optical Co. Ltd., Tokyo, Japan) with distal cap attachment and the Lucera system video endoscopy processor (Olympus CV-260). This equipment provides the advanced features of adaptive index of haemoglobin (IHb) colour and structural enhancement, which emphasises subtle chromatic alterations in the mucous membrane. We visualised coil-like capillary loops of normal antral mucosa (white arrows, upper left), a clear demarcation line between cancerous and noncancerous mucosa (black arrows), and a leash of irregular vessels suggestive of neovascularisation and angiogenesis secondary to dysplasia (grey arrows). Targeted biopsies were obtained and histological examination revealed high-grade dysplasia.