


Figure 1 A 47-year-old man presented with a 2-week history of mid-abdominal pain. He denied having any fever, chills, nausea, vomiting, or diarrhea. Computed tomography of the abdomen revealed numerous air-filled cysts involving the splenic flexure of the colon (arrow), a finding consistent with pneumatosis coli. A colonoscopy examination was then performed, which revealed sigmoid diverticulosis.






Figure **2** Multiple large, intramural, cyst-like structures were seen at the level of the splenic flexure at colonoscopy.

T. E. Yusuf¹, S. C. Abraham², S. W. Trenkner³

- ¹ Division of Gastroenterology and Hepatology, Mayo Clinic, Rochester, Minnesota, USA
- ² Division of Anatomic Pathology, Mayo Clinic, Rochester, Minnesota, USA
- ³ Department of Radiology, Mayo Clinic, Rochester, Minnesota, USA

Corresponding Author

T. E. Yusuf M. D.

Advanced Endoscopy Unit Division of Gastroenterology and Hepatology Blake 4 Massachusetts General Hospital 55 Fruit Street Boston Massachusetts 02114 USA

Fax: +1-617-724-5997 E-mail: tyusuf@partners.org




Figure 3 Histopathologic examination of biopsy specimens revealed cystic airfilled spaces within the submucosa which were partially lined by clusters of foreignbody macrophages (hematoxylin & eosin stain, magnification × 10). A few weeks later the patient developed a perforated diverticulitis, which was managed medically; sigmoidectomy is planned.