

There has been a resurgence of intestinal tuberculosis during the last decade following the increase of immigrants to Western countries [1,2]. The abdominal symptoms of intestinal tuberculosis are nonspecific, consisting mainly of abdominal pain, weight loss, anorexia, and fever [2,3]. The characteristic endoscopic features of colonic tuberculosis include transversely oriented ulcer, nodules, deformed ileocecal valve, stricture, erosions, and aphthous ulcers [1–3]. However, in patients without abdominal symptoms or pulmonary infection, endoscopists face the challenge of distinguishing between intestinal tuberculosis and malignancy or Crohn's disease. We report here a case of colonic tuberculosis found incidentally during a health check-up, and which was diagnosed by typical endoscopic features and histological and microbiological evidence.

A 38-year-old, previously healthy man visited our institution for a scheduled health check-up. His medical history and physical examination were unremarkable. He had a history of travel to China 3 months before coming to our hospital. Chest radiograph showed no active pulmonary lesions. Laboratory data were within reference range except for a triglyceride level of 381 mg/dL (normal range 50–130 mg/dL). Tumor markers including CEA, CA 125 and CA 19–9 were normal. Ziehl–Neelsen stain and culture of sputum were negative for *Mycobacterium tuberculosis*.

Colonoscopy revealed a transversely oriented ulcer in the cecum, with steep edges and surrounding flared nodules (Figure 1). The terminal ileum appeared normal. Histological examination of the biopsy specimens demonstrated well-formed granulomas with caseous necrosis and Langhan's giant cells (Figure 2). Culture of biopsy specimens revealed positivity for *M. tuberculosis*, as did the result of polymerase chain reaction for DNA of *M. tuberculosis*. A diagnosis of colonic tuberculosis was made. The patient received




Figure 1 Colonoscopic view showing a transversely oriented ulcer in the cecum with steep edges and flared surrounding nodules. Note its location opposite the ileocecal orifice (arrow).




Figure 2 Histological view showing well-formed granulomas with Langhan's giant cells (arrows) and caseous necrosis (n), surrounded by a prominent rim of lymphocytes (hematoxylin and eosin; original magnification $\times 100$).

antitubercular therapy for 9 months with an uneventful clinical course. He remained asymptomatic over the ensuing 2 years of follow-up.

Endoscopy_UCTN_Code_CCL_1AD_2AC

C. C. Yang, C. H. Chen, S. L. Yan

Division of Gastroenterology, Department of Internal Medicine, Show-Chwan Memorial Hospital, Taiwan, Republic of China.

References

- 1 Sato S, Yao K, Yao T et al. Colonoscopy in the diagnosis of intestinal tuberculosis in asymptomatic patients. *Gastrointest Endosc* 2004; 59: 362–368
- 2 Alvares JF, Devarbhavi H, Makhija P et al. Clinical, colonoscopic, and histological profile of colonic tuberculosis in a tertiary hospital. *Endoscopy* 2005; 37: 351–356
- 3 Misra SP, Misra V, Dwivedi M, Gupta SC. Colonic tuberculosis: clinical features, endoscopic appearance and management. *J Gastroenterol Hepatol* 1999; 14: 723–729

Corresponding author

S. L. Yan, MD

Division of Gastroenterology
Department of Internal Medicine
Show-Chwan Memorial Hospital
No 542, Sec 1, Chung-Shang Rd
Changhua 500
Taiwan
Republic of China
Fax: +886-4-7233190
Email: yslscmc@yahoo.com