



**Figure 1** A 55-year-old woman with systemic lupus erythematosus, who had been followed up regularly for 7 years, visited our hospital complaining of chronic abdominal fullness, constipation, and stringy stools. She had a history of a total hysterectomy for uterine prolapse at the age of 38. At colonoscopy we found a 3-cm-long tubular structure, with normal-appearing mucosa, in the cecum; histopathological examination showed colonic mucosa with mild inflammation. A diagnosis of inverted appendix was made.



**Figure 2** Virtual colonoscopy performed immediately after colonoscopy revealed a three-dimensional tubular structure, similar to that seen on colonoscopy. Inverted appendix should be considered in the differential diagnosis of cecal polyps.

**A.-L. Chou<sup>1</sup>, C.-W. Lin<sup>2</sup>, K.-C. Tseng<sup>1</sup>**

<sup>1</sup> Department of Internal Medicine, Buddhist Dalin Tzu Chi General Hospital, Chiayi, Taiwan

<sup>2</sup> Department of Radiology, Buddhist Dalin Tzu Chi General Hospital, Chiayi, Taiwan

Endoscopy\_UCTN\_Code\_CCL\_1AD\_2AC

Endoscopy\_UCTN\_Code\_CCL\_1AD\_2AJ

## Corresponding Author

**K.-C. Tseng, M.D.**

Department of Internal Medicine  
Buddhist Dalin Tzu Chi General Hospital  
2 Min Sheng Road  
Dalin  
Chiayi  
Taiwan

Fax: +886-5-2648006

E-mail: dennis0.tseng@msa.hinet.net