

## Iron-induced esophageal ulceration

Esophageal inflammation or ulceration is a well recognized side-effect of some oral medications, especially nonsteroidal anti-inflammatory or potassium chloride tablets. A similar injury can be induced by iron but endoscopically detected lesions are rarely reported [1,2].

An 82-year-old woman presented a sudden complaint of dysphagia after ingestion of a ferrous sulphate tablet. She had chronic esophageal reflux and Raynaud's syndrome. Her physical examination was normal.

An upper endoscopy revealed a black-brown, ulcerated and necrotic lesion, just below the upper esophageal sphincter, 2 cm in length, well circumscribed and associated with a slight stenosis. Below the lesion the esophageal lumen was distended and atonic, with candidiasis and a grade D peptic esophagitis (◀ Fig. 1, ▶ Video 1).

Histological evaluation revealed ulcerated esophagitis, with acute inflammatory granulation tissue containing abundant brown crystalline nonbirefringent material, confirmed as iron by Perl's staining (▶ Fig. 2, 3). Other stainings for brown pigments and infections were negative.

The patient's iron tablets were stopped and a high dose proton pump inhibitor was prescribed. The dysphagia was relieved and a repeat upper endoscopy revealed progressive resolution of the necrosis and stenosis (▶ Fig. 4, ▶ Video 2). Iron-induced esophageal erosions or ulcerations are seldom described and the differential diagnoses might be striking. Tablet debris seen at endoscopy is diagnostic but rarely detected. Histological examination reveals a typical brown crystalline material positive with Perl's staining [1,3,4]. The pathogenesis is believed to be a high local iron saturation causing a concentration-dependent absorption, resulting in the formation of reactive oxygen metabolites and mucosal injury [3]. Anatomical factors such as motility disorders or previous inflammation might predispose to the iron injury [3,4].

### Video 1

Upper endoscopy at day 1 revealed a black-brown, ulcerated and necrotic lesion, just below the upper esophageal sphincter and a distended atonic esophagus.

Treatment requires stopping the offending drug or, if iron is still necessary, changing to a liquid preparation [3,5]. Iron-induced lesions are not endoscopically specific and histological investigation is essential to the diagnosis.

Endoscopy\_UCTN\_Code\_CCL\_1AB\_2AC\_3AZ

**M. Areia<sup>1</sup>, R. Gradiz<sup>1</sup>, P. Souto<sup>1</sup>, E. Camacho<sup>1</sup>, M. R. Silva<sup>2</sup>, N. Almeida<sup>1</sup>, A. Rosa<sup>1</sup>, M. F. Xavier da Cunha<sup>2</sup>, M. C. Leitão<sup>1</sup>**

<sup>1</sup> Department of Gastroenterology, Coimbra University Hospital, Coimbra, Portugal

<sup>2</sup> Department of Pathology, Coimbra University Hospital, Coimbra, Portugal

### References

- 1 Abbarah TR, Fredell JE, Ellenz GB. Ulceration by oral ferrous sulfate. *JAMA* 1976; 236: 2320
- 2 Zhang ST, Wong WM, Hu WH et al. Esophageal injury as a result of ingestion of iron tablets. *J Gastroenterol Hepatol* 2003; 18: 466–467
- 3 Abraham SC, Yardley JH, Wu TT. Erosive injury to the upper gastrointestinal tract in patients receiving iron medication: an underrecognized entity. *Am J Surg Pathol* 1999; 23: 1241–1247
- 4 Eckstein RP, Symons P. Iron tablets cause histopathologically distinctive lesions in mucosal biopsies of the stomach and esophagus. *Pathology* 1996; 28: 142–145
- 5 Haig A, Driman DK. Iron-induced mucosal injury to the upper gastrointestinal tract. *Histopathology* 2006; 48: 808–812

### Bibliography

DOI 10.1055/s-2007-966820

Endoscopy 2007; 39: E326

© Georg Thieme Verlag KG Stuttgart · New York · ISSN 0013-726X


### Corresponding author

**M. Areia, MD**


Department of Gastroenterology  
Coimbra University Hospital  
3000-075 Coimbra  
Portugal  
Fax: +351-239-701517  
miguel.areia@netcabo.pt

### Video 2


Upper endoscopy at day 8 showing the progressive resolution of the lesion with a circumferential brown-pigmented lesion with no necrosis or stenosis.



**Fig. 1** Upper endoscopy at day 1 revealed a black-brown, ulcerated and necrotic lesion, just below the upper esophageal sphincter.



**Fig. 2** Granulation tissue from the area with ulcerated esophagitis containing abundant brown crystalline material (hematoxylin and eosin, ×400).



**Fig. 3** Positive blue staining of the crystalline material with Perl's iron stain (×400).



**Fig. 4** Upper endoscopy at day 8 showing the progressive resolution of the lesion, with a brown pigmentation and no necrosis.