Invasive cancer arising in a colonic inverted hyperplastic polyp



Fig. 1 Colonoscopy showed a whitish, flat, elevated polyp, 10 mm in diameter, with a dimple on the top in the ascending colon.

A 64-year-old man underwent colonoscopy for the investigation of lower abdominal pain. Colonoscopy showed a whitish, flat, elevated polyp, 10 mm in diameter, in the ascending colon (**S Fig. 1**). Endoscopic mucosal resection (EMR) was carried out, and the resected specimen histologically revealed both invasive cancer and serrated adenomatous glands coexisting in an inverted hyperplastic polyp (IHP) (**Fig. 2a-c**). A follow-up colonoscopy 6 months later revealed an irregularly elevated normal mucosa with fold convergence adjacent to the EMR scar (**Fig. 3**). Although endoscopic features suggested a recurrent cancer, endoscopic biopsies unfortunately failed to identify recurrent cancer histologically. Surprisingly, a second follow-up colonoscopy only 3 months later showed an advanced colon cancer (> Fig. 4). Endoscopic biopsies showed poorly differentiated adenocarcinoma, and therefore surgical resection was carried out. Histologically, the recurrent cancer was a subserosally invasive, poorly differentiated adenocarcinoma with vascular invasion and nodal involvement, and it was classified as stage III (T3N2M0).

This IHP was 10 mm in diameter, which is a so-called large hyperplastic polyp, and interestingly it demonstrated a characteristic endoscopic and histologic feature of IHP [1]. IHP is a variant form of hyperplastic polyp with a characteristic histologic



feature of epithelial misplacement or inversion of the epithelium into the submucosa [2]. Our case was unique and the histology detected the presence of not only an invasive, poorly differentiated carcinoma but also serrated adenomatous glands surrounded by hyperplastic glands. To date, three cases of IHP associated with adenoma have been reported [3]; however this is the first case report of both of invasive cancer and serrated adenoma arising in an IHP. Recently, serrated polyps have been histologically defined as polyps that demonstrate a serrated or "sawtoothed" appearance, and include hyperplastic polyps, serrated adenoma, and



Fig. 3 At the first follow-up colonoscopy, an irregularly elevated normal mucosa with fold convergence mimicking a submucosal tumor was detected at the resection site. Endoscopic features suggested a submucosal mass probably created by a recurrent cancer.



Fig. 4 An ulcerative lesion, suggesting an advanced colon cancer, was detected at the site of the resection scar.

polyps with combined histology [4]. Carcinoma associated with serrated adenoma has been reported to be a distinct type of colorectal neoplasm, accounting for 5.8% to 7.5% of all colorectal carcinoma cases and up to 17.5% of proximal colon cancers [5]. Based on the concept of the "serrated polyp neoplasia pathway", the serrated adenoma in this IHP could be considered to originate from the hyperplastic glands, as it was surrounded by hyperplastic glands especially in the inverted part, and it would subsequently replace the surrounding hyperplastic glands and then develop into invasive carcinoma with undifferentiated histology.

Endoscopy_UCTN_Code_CCL_1AD_2AB

K. Fu¹, T. Fujii², H. Kuwayama³, T. Ishikawa⁴, Y. Ueda⁵, T. Fujimori⁶

- ¹ Department of Coloproctology, Tokatsu-Tsujinaka Hospital, Chiba, Japan
- ² TF Clinic, Ginza, Tokyo, Japan
- ³ Department of Gastroenterology and Hepatology, Koshigaya Hospital, Dokkyo Medical University, Koshigaya, Saitama, Japan
- ⁴ Department of Radiology, Dokkyo Medical University, Tochigi, Japan
- ⁵ Department of Pathology, Koshigaya Hospital, Dokkyo Medical University, Koshigaya, Saitama, Japan
- ⁶ Department of Surgical and Molecular Pathology, Dokkyo Medical University, Tochigi, Japan

References

- 1 *Fu KI, Matsuda T, Saito Y et al.* An inverted hyperplastic polyp with a characteristic colonoscopic appearance. Endoscopy 2006; 38 Suppl 2: E55
- 2 Sobin LH. Inverted hyperplastic polyps of the colon. Am J Surg Pathol 1985; 9: 265 – 272
- 3 *Kuribayashi K, Ishii T, Ishidate T et al.* Two cases of inverted hyperplastic polyps of the colon and association with adenoma. Eur J Gastroenterol Hepatol 2004; 16: 107–112
- 4 Hawkins NJ, Bariol C, Ward RL. The serrated neoplasia pathway. Pathology 2002; 34: 548-555
- 5 *Mäkinen MJ*. Colorectal serrated adenocarcinoma. Histopathology 2007; 50: 131 – 150

Bibliography

DOI 10.1055/s-2007-995733 Endoscopy 2010; 42: E29–E30 © Georg Thieme Verlag KG Stuttgart · New York · ISSN 0013-726X

Corresponding author

K. Fu, MD, PhD Department of Coloproctology Tokatsu-Tsujinaka Hospital Chiba, 270-1168 Japan Fax: +81-4-71849854 fukuangi@hotmail.com