Case Report

Abdominal compartment syndrome following abdominoplasty: A case report and review

Arash Izadpanah, Ali Izadpanah¹, Mihiran Karunanayake¹, Christian Petropolis, Dan L. Deckelbaum², Mario Luc¹

Division of Plastic and Reconstructive Surgery, University of Manitoba, Health Sciences Centre, Winnipeg, Manitoba, ¹Division of Plastic and Reconstructive Surgery, ²Division of Trauma, McGill University, McGill University Health Centre, Montreal, Quebec, Canada

Address for correspondence: Dr. Ali Izadpanah, Division of Plastic and Reconstructive Surgery, McGill University Health Centre, 760 Upper Lansdowne Avenue, Westmount, Quebec H3Y 1J8, Canada. E-mail: ali.izadpanah@mail.mcgill.ca

ABSTRACT

Abdominoplasty is among the most commonly performed aesthetic procedures in plastic surgery. Despite high complication rate, abdominal contouring procedures are expected to rise in popularity with the advent of bariatric surgery. Patients with a history of gastric bypass surgery have an elevated incidence of small bowel obstruction from internal herniation, which is associated with non-specific upper abdominal pain, nausea, and a decrease in appetite. Internal hernias, when subjected to elevated intra-abdominal pressures, have a high-risk of developing ischemic bowel. We present a case report of patient with previous laparoscopic Roux-en-y gastric bypass who developed acute ischemic bowel leading to abdominal compartment syndrome following abdominoplasty. To the best of our knowledge, this is the first reported case in the literature. We herein emphasise on the subtle symptoms and signs that warrant further investigations in prospective patients for an abdominal contouring procedure with a prior history of gastric bypass surgery.

KEY WORDS

Abdminoplasty; abdominal compartment syndrome; bariatric surgery

INTRODUCTION

ccording to the American Society of Plastic Surgeons, abdominoplasty is the 5th most common cosmetic surgery in the United States.^[1] Patients often request abdominal wall contouring procedures due to excessive skin and fat following pregnancy, excessive weight loss and fat maldistribution, which

Access this article online				
Quick Response Code:	Website: www.ijps.org			
	DOI: 10.4103/0970-0358.138978			

can lead to difficulties with exercise, sexual activities, fitting into clothing and hygiene.^[2] Despite its popularity, abdominoplasty is not devoid of complications. The complications rates remain high, ranging from 34 to 37.4%.^[3-5] The most common reported complications are seroma formation followed by wound complications and skin flap necrosis.^[6-8] Less common complications include hematoma formation, umbilical necrosis, wound dehiscence, contour deformities, and abnormal scarring.^[9] In addition, there are only a few reports of increased intra-abdominal pressures in patients who had recently undergone an abdominoplasty.^[10-12] Although very small, there is a potential risk of bowel ischemia following abdominoplasty in patients with pervious history of intra-abdominal surgeries.

To best of our knowledge, there have been no previous reports of abdominal compartment syndrome following an abdominoplasty in a patient with history of gastric bypass surgery. As such, surgeons should be aware of increased risk of ischaemic bowel that can occur in post-bariatric patients who undergo abdominal contouring procedures.

CASE REPORT

A 30-year-old female, G₀P₀A₀ non-smoking nurse underwent an abdominoplasty by the senior author. Her past medical history was remarkable for a laparoscopic Roux-en-Y gastric bypass (LRYGB) performed 1-year prior to current presentation. Her initial weight prior to LRYGB was 108 kg with body mass index (BMI) of 42. She subsequently lost over 120 pounds and developed a significant abdominal pannus. Her weight has been stable over 6 months prior to current presentation with her new BMI at 21. She underwent bilateral breast augmentation and an abdominoplasty, without fascia plication and with the use of a standard abdominal binder. The patient was discharged home on the postoperative day 1 in stable conditions. This is the common practice for all uncomplicated abdominoplasties in our institution.

On post-operative day 2, she developed persistent nausea, low urine output, generalised weakness and severe abdominal pain. She presented to the emergency department and was found to be febrile, tachypnoeic and tachycardic. Her abdominal examination revealed significant tenderness and guarding. Computed tomography revealed significant intra-abdominal free fluid and suggestion of pneumatosis intestinalis. While being aggressively resuscitated, the patient became haemodynamically unstable unresponsive to fluids, developed respiratory failure, rising lactate levels (6.7 mmol/L), elevated peak ventilator pressures and was urgently brought into the operating room with the presumptive diagnosis of mesenteric ischemia and abdominal compartment syndrome.

Patient underwent a laparotomy procedure. Upon entry of the peritoneal cavity, her haemodynamics and respiratory status improved immediately. The alimentary limb of the previous Roux-en-Y bypass was completely necrotic all the way from the pouch to the level of the jejunojejunostomy. The volvulised alimentary as it passes through the transverse mesocolon was excised. A nasogastric tube was inserted intraoperatively to decompress the pouch. 48 h later, a side-to-side gastrogastrostomy between the gastric pouch and the gastric remnant was performed. Her post-operative course was complicated by pneumonia and a localised laparotomy wound infection. Both were treated appropriately with intravenous antibiotics and local wound debridement. She was initially discharged from the hospital on the post-operative day twelve with a vacuum assisted device applied to her abdominal defect. The final closure of this defect was carried out 20 days following discharged with a split thickness skin graft.

DISCUSSION

Abdominoplasty is amongst the most commonly requested aesthetic procedures in United States. The demand for abdominoplasty is expected to rise with the advent of bariatric procedures. It is therefore imperative that plastic surgeons be aware of all complications that can occur following an abdominoplasty in patients who have previously undergone a bariatric procedure.

The Roux-en-Y gastric bypass (RYGB) is the most common and the standard of care for surgical treatment of obesity.^[13] The majority of weight loss has been shown to occur within the first 12-18 months through restricted intake, and/or malabsorption.^[14] Patients requesting body contouring surgery following bariatric surgery can often have chronic nutritional deficiencies, predisposing them to a risk of impaired wound healing.^[15] A stable weight over 6 months prior to an abdominoplasty has been shown to be associated with less wound healing complications and better aesthetic outcomes.^[15] Our patient fit this category and therefore there were no concerns about her nutritional status preoperatively.

The months prior to her surgery, she was however complaining of occasional recurrent abdominal discomfort related to food intake. These symptoms could possibly be attributed to the herniation of the alimentary limb of the gastric bypass through the transverse mesocolon as discovered during the exploratory laparotomy. Herniation leading to volvulous subsequent to a RYGB in our patient could have been attributed to the post-surgical adhesions and presence of a mesenteric defect at the transverse mesocolon.^[16] Performing a simple non-invasive abdominal ultrasound in patients with pervious history of gastric bypass could potentially help in diagnosing internal and external hernias and should be routine in this patient population.

Internal herniation of the small bowel has reported incidence ranging from 1.8% to 4.5% in patients who have had a LRYGB.^[17-20] In addition, pregnancy has been demonstrated to increase the likelihood of an internal hernia in patients with a RYGB.^[21-24] Herniation occurs at the jejunojejunostomy site through the transverse mesocolon as seen in our patient, or through Peterson's defect [Figure 1].^[25] The initial symptoms and signs in our patient were non-specific and included epigastric or periumbilical pain, decreased appetite and nausea, which is in keeping with previous reports.^[25]

Imaging studies are not always diagnostic. Wang *et al.* concluded that unrelenting epigastric pain in patients



Figure 1: Potential spaces for internal herniation following Roux-en-Y gastric bypass surgery. (a) Petersonæs hernia, through the posterior aspect of the mesentery of the Roux limb. (b) Transverse mesocolon, as witnessed in our patient. (c) Jejunomesenteric hernia through the potential space from the jejunojenostomy

with a history of RYGB surgery should raise concern for small bowel obstruction and exploratory laparotomy or diagnostic laparoscopy should be considered.^[21] Failing to investigate these patients for internal herniation places them at risk of complications such as ischemic bowel when subjected to elevations in intra-abdominal pressure.

Charles *et al.* in their have reported a patient in her late second trimester having undergone a RYGB 6 months prior to the presentation, that subsequently developed ischemic bowel secondary to an internal herniation of the afferent limb from the raised intra-abdominal pressure due to pregnancy.^[24] The patient survived but delivered a nonviable foetus in the immediate postoperative period.

Although there have been reports of elevated intraabdominal pressures post-abdominoplasty, there have been no reported cases of ischaemic bowel^[10,26] [Table 1]. A delayed small bowel obstruction 2 years following an abdominoplasty in a massive weight loss patient with gastric bypass has been previously reported.^[27] The patient suffered from an ischaemic bowel obstruction secondary to migration of a gastric band's port chamber of in the anterior abdominal wall.

Abdominal plication and use of binders has been demonstrated to be associated with an increase in abdominal pressure after abdominoplasty.^[11] Thus, in the absence of plication of the rectus abdominis, we suspect that the ischemia may have resulted primarily from strangulation of an internal hernia due to elevated intra-abdominal pressure from the standard abdominal binder. This was further exacerbated due to the presence of possible hypotension secondary to nausea and lower fluid intake and by excessive narcotic intake in immediate post-operative course. The resultant inflammation and oedema of the necrotic bowel further

Publication	Authors	Complications	Treatment	Outcome	
2011	Izadpanah et al.	1 case of ACS	Laparotomy decompression	Satisfactory immediately post- operatively	
2007	Shen <i>et al</i> .*	3 cases of ACS	Two patients opted for laparotomy decompression	Satisfactory in 2 years follow-up	
2006	Graça Neto <i>et al.</i>	0 cases of ACS Two patients developed mild dyspnoea with mildly elevated abdominal compartment pressures	No treatment was necessary	Not mentioned	

Table 1: Reported cases of increased abdominal pressure following abdominoplasty

*Only the abstract was accessible; ACS: Abdominal compartment syndrome

elevated the intra-abdominal pressures causing the suspected abdominal compartment syndrome.

CONCLUSION

In conclusion, with the rise in demand for abdominal contouring procedures from patients who have undergone bariatric surgery, plastic surgeons should be aware of the subtle clinical symptoms and signs, such as abdominal pain, that suggest underlying propensity for internal herniation, which could eventually lead to small bowel obstruction or ischaemia. Furthermore, abdominal compartment syndrome, although rare following abdominoplasty, can be a product of ischemia of an internal herniation of a limb from a previous LRYGP when subjected to elevations in intra-abdominal pressure, leading to patient morbidity and increased health care costs.

REFERENCES

- American Society of Plastic Surgeons. Available from: http:// www.plasticsurgery.org/News-and-Resources/Statistics.html. [Last accessed on Jul 2011, cited on Jul 2011].
- Breiting LB, Lock-Andersen J, Matzen SH. Increased morbidity in patients undergoing abdominoplasty after laparoscopic gastric bypass. Dan Med Bull 2011;58:A4251.
- Hensel JM, Lehman JA Jr, Tantri MP, Parker MG, Wagner DS, Topham NS. An outcomes analysis and satisfaction survey of 199 consecutive abdominoplasties. Ann Plast Surg 2001;46:357-63.
- Floros C, Davis PK. Complications and long-term results following abdominoplasty: A retrospective study. Br J Plast Surg 1991;44:190-4.
- Neaman KC, Hansen JE. Analysis of complications from abdominoplasty: A review of 206 cases at a university hospital. Ann Plast Surg 2007;58:292-8.
- Hafezi F, Nouhi AH. Abdominoplasty and seroma. Ann Plast Surg 2002;48:109-10.
- van Uchelen JH, Werker PM, Kon M. Complications of abdominoplasty in 86 patients. Plast Reconstr Surg 2001;107:1869-73.
- Chaouat M, Levan P, Lalanne B, Buisson T, Nicolau P, Mimoun M. Abdominal dermolipectomies: Early postoperative complications and long-term unfavorable results. Plast Reconstr Surg 2000;106:1614-8.
- Momeni A, Heier M, Bannasch H, Stark GB. Complications in abdominoplasty: A risk factor analysis. J Plast Reconstr Aesthet Surg 2009;62:1250-4.
- Graça Neto L, Araújo LR, Rudy MR, Auersvald LA, Graf R. Intraabdominal pressure in abdominoplasty patients. Aesthetic Plast Surg 2006;30:655-8.

- Huang GJ, Bajaj AK, Gupta S, Petersen F, Miles DA. Increased intraabdominal pressure in abdominoplasty: Delineation of risk factors. Plast Reconstr Surg 2007;119:1319-25.
- Shen GX, Gu B, Xie F, Lu K, Wang HY, Zheng DN, *et al.* Three case reports of abdominal compartment syndrome after full abdominoplasty. Zhonghua Zheng Xing Wai Ke Za Zhi 2007;23:226-8.
- Tice JA, Karliner L, Walsh J, Petersen AJ, Feldman MD. Gastric banding or bypass? A systematic review comparing the two most popular bariatric procedures. Am J Med 2008;121:885-93.
- Buchwald H, Avidor Y, Braunwald E, Jensen MD, Pories W, Fahrbach K, et al. Bariatric surgery: A systematic review and meta-analysis. JAMA 2004;292:1724-37.
- Agha-Mohammadi S, Hurwitz DJ. Nutritional deficiency of postbariatric surgery body contouring patients: What every plastic surgeon should know. Plast Reconstr Surg 2008;122:604-13.
- 16. Macgregor AM. Small bowel obstruction following gastric bypass. Obes Surg 1992;2:333-339.
- Champion JK, Williams M. Small bowel obstruction and internal hernias after laparoscopic Roux-en-Y gastric bypass. Obes Surg 2003;13:596-600.
- Garza E Jr, Kuhn J, Arnold D, Nicholson W, Reddy S, McCarty T. Internal hernias after laparoscopic Roux-en-Y gastric bypass. Am J Surg 2004;188:796-800.
- Hwang RF, Swartz DE, Felix EL. Causes of small bowel obstruction after laparoscopic gastric bypass. Surg Endosc 2004;18:1631-5.
- Higa KD, Ho T, Boone KB. Internal hernias after laparoscopic Roux-en-Y gastric bypass: Incidence, treatment and prevention. Obes Surg 2003;13:350-4.
- Wang CB, Hsieh CC, Chen CH, Lin YH, Lee CY, Tseng CJ. Strangulation of upper jejunum in subsequent pregnancy following gastric bypass surgery. Taiwan J Obstet Gynecol 2007;46:267-71.
- Moore KA, Ouyang DW, Whang EE. Maternal and fetal deaths after gastric bypass surgery for morbid obesity. N Engl J Med 2004;351:721-2.
- Kakarla N, Dailey C, Marino T, Shikora SA, Chelmow D. Pregnancy after gastric bypass surgery and internal hernia formation. Obstet Gynecol 2005;105:1195-8.
- Charles A, Domingo S, Goldfadden A, Fader J, Lampmann R, Mazzeo R. Small bowel ischemia after Roux-en-Y gastric bypass complicated by pregnancy: A case report. Am Surg 2005;71:231-4.
- Eckhauser A, Torquati A, Youssef Y, Kaiser JL, Richards WO. Internal hernia: Postoperative complication of roux-en-Y gastric bypass surgery. Am Surg 2006;72:581-4; discussion 584.
- Talisman R, Kaplan B, Haik J, Aronov S, Shraga A, Orenstein A. Measuring alterations in intra-abdominal pressure during abdominoplasty as a predictive value for possible postoperative complications. Aesthetic Plast Surg 2002;26:189-92.
- 27. SarkarS, SmithD, ScottMH.Arare complication of abdominoplasty after bariatric surgery. Ann Plast Surg 2010;64:7-8.

How to cite this article: Izadpanah A, Izadpanah A, Karunanayake M, Petropolis C, Deckelbaum DL, Luc M. Abdominal compartment syndrome following abdominoplasty: A case report and review. Indian J Plast Surg 2014;47:263-6.

Source of Support: Nil, Conflict of Interest: None declared.