

Alopecia due to cancer

Sir,

Cancer is the group of disease presenting with uncontrolled cell growth that lead to the pathologies of the patients. Interestingly, cancer can be the cause of alopecia. This is a topic to be considered in trichology. Here, the author will presents and discuss on some important malignancies. The problem is frequently mentioned in the hematological malignancies. Of several hematological cancers, the alopecia is mentioned as an important clinical manifestation of mycosis

fungoides and Sézary syndrome.^[1] Bi *et al.* reported that alopecia could be observed in 2.5% of these patients.^[2] Furthermore, Bi *et al.* observed that alopecia could be detectable within mycosis fungoides lesions in 66%.^[2] The possible concomitant skin lesions in patients include disseminated follicular erythematous papules, and comedones and cysts.^[3] The cutaneous lymphoma should be usually included into the differential diagnosis of alopecia, especially for alopecia at scalp.^[1,4-6] An important condition that is very similar to the cutaneous lymphoma related alopecia is secondary syphilis related alopecia in

human immunodeficiency virus infected patients.^[7] In addition to lymphoma, alopecia is also mentioned in plasmacytoma.^[8] Alopecia is an important component in paraneoplastic syndrome due to plasmacytoma.^[8,9] In addition to hematologic malignancies, the solid cancers can also cause alopecia. Alopecia might be induced by tumor metastasis to the scalp.^[10] The common underlying source of metastasis is breast cancer in female.^[10] The other uncommon solid tumor that can cause alopecia is adenoma.^[11] The mechanism is due to the hyperandrogenemia.^[11]

However, the more well-known cancer-related alopecia is the side-effect of cancer therapy. It is no doubt that chemotherapy can induce alopecia. This problem can be seen in three-fifths of the patients undergoing chemotherapy.^[12] The pathological changes of hair include trichorrhexis, fragmentation, decreased hair diameter, and hair shaft depigmentation.^[12] Focusing on the natural history, the problem usually occurs after the 1st or the 2nd cycle of administration of therapeutic regimen (about 1 week to 2 months after the start of cancer therapy).^[12] More numbers of used chemotherapeutic agents result in more severe alopecia.^[12] For reducing the problem, immunomodulator drugs, both modern and herbal regiment seems to be useful in the management of this problem.^[12] Furthermore, the poor nutritional status increases the risk of alopecia, hence the improvement of nutrition for patients receiving chemotherapy can be a good preventive method for this problem.^[13]

Somsri Wiwanitkit¹, Viroj Wiwanitkit^{1,2,3,4}

¹Wiwanitkit House, Bangkhae, Bangkok, Thailand,

²Visiting University, Hainan Medical University, China,

³Joesph Ayo Babalola University, Nigeria,

⁴Faculty of Medicine, University of Nis, Serbia.

E-mail: somsriwiwan@hotmail.com

REFERENCES

1. Monopoli A, Annessi G, Lombardo GA, Baliva G, Girolomoni G. Purely follicular mycosis fungoides without mucinosis: Report of two cases with review of the literature. *J Am Acad Dermatol* 2003;48:448-52.
2. Bi MY, Cohen PR, Robinson FW, Gray JM. Alopecia

syphilitica-report of a patient with secondary syphilis presenting as moth-eaten alopecia and a review of its common mimickers. *Dermatol Online J* 2009;15:6.

3. Bi MY, Curry JL, Christiano AM, Hordinsky MK, Norris DA, Price VH, *et al*. The spectrum of hair loss in patients with mycosis fungoides and Sézary syndrome. *J Am Acad Dermatol* 2011;64:53-63.
4. Colovic N, Jurisic V, Terzic T, Atkinson HD, Colovic M. Immunochemotherapy for Bcl-2 and MUM-negative aggressive primary cutaneous B-cell non-Hodgkin's lymphoma. *Arch Dermatol Res* 2009;301:689-92.
5. Török L, Gurbity TP, Kirschner A, Krenács L. Panniculitis-like T-cell lymphoma clinically manifested as alopecia. *Br J Dermatol* 2002;147:785-8.
6. Török L, Gurbity PT, Kirschner A, Krenács L. Panniculitis-like lymphoma clinically manifesting as alopecia. *Orv Hetil* 2002;143:607-9.
7. Baniandrés Rodríguez O, Nieto Perea O, Moya Alonso L, Carrillo Gijón R, Harto Castaño A. Nodular secondary syphilis in a HIV patient mimicking cutaneous lymphoma. *An Med Interna* 2004;21:241-3.
8. Weichenthal M, Stemm AV, Ramsauer J, Mensing H, Feller AC, Meigel W. POEMS syndrome: Cicatricial alopecia as an unusual cutaneous manifestation associated with an underlying plasmacytoma. *J Am Acad Dermatol* 1999;40:808-12.
9. Wheeler GE, Barrows GH. Alopecia universalis. A manifestation of occult amyloidosis and multiple myeloma. *Arch Dermatol* 1981;117:815-6.
10. Chen W, Yang CC, Todorova A, Al Khuzaei S, Chiu HC, Worret WI, *et al*. Hair loss in elderly women. *Eur J Dermatol* 2010;20:145-51.
11. Chen W, Chen GY, Tsai SJ, Wang PW, Sheu HM, Shen YS, *et al*. Mild cutaneous manifestation in two young women with extraordinary hyperandrogenemia. *Dermatology* 2005;210:49-52.
12. Pai GS, Vimala AM, Dinesh M. Occurrence and severity of alopecia in patients on combination chemotherapy. *Indian J Cancer* 2000;37:95-104.
13. Arrieta O, Michel Ortega RM, Villanueva-Rodríguez G, Serna-Thomé MG, Flores-Estrada D, Diaz-Romero C, *et al*. Association of nutritional status and serum albumin levels with development of toxicity in patients with advanced non-small cell lung cancer treated with paclitaxel-cisplatin chemotherapy: A prospective study. *BMC Cancer* 2010;10:50.

Access this article online

Quick Response Code:



Website:
www.ijmpo.org

DOI:
10.4103/0971-5851.116228