## **CASE REPORT**



# Giant cell tumor of the clivus with presence of epithelioid histiocytes

Amit Agrawal, Rajsekhar Gali<sup>1</sup>, Vissa Shanthi<sup>2</sup>, Baddukonda Appala Ramakrishna<sup>2</sup>, Kuppili Venkata Murali Mohan<sup>2</sup>

Department of Neurosurgery, Narayana Medical College Hospital, Chinthareddypalem, Departments of 1Oral and Maxillofacial Surgery, <sup>2</sup>Department of Pathology, Narayana Dental College Hospital, Chinthareddypalem, Nellore, Andhra Pradesh, India

#### ABSTRACT

Giant cell tumor (GCT) is a benign neoplasm but locally aggressive tumor that uncommonly involves the skull bone. We report a case of a 62-year-old male presented with increasing headache and diplopia. Investigations were suggestive of an expanding mass lesion of the clivus. Histopathology was suggestive of diagnosed with GCT with abundant histiocytes.

Key words: Clivus, giant cell tumor, histiocytes, skull

## Introduction

Giant cell tumor (GCT) is a benign neoplasm but locally aggressive tumor that uncommonly involves the skull bone (preferentially, the sphenoid and the temporal bones),<sup>[1,2]</sup> with ooccasional instances of orbital, calvarial, and occipital bones involvement.<sup>[3]</sup> In the present article, we report a case of GCT of the clivus and review the relevant literature.

### **Case Report**

A 62-year-old male underwent transphenoidal biopsy of the sellar mass lesion and it was diagnosed as GCT. Now, he presented with an increasing in headache and diplopia of 3 months duration. He was a known diabetic controlled with medication. There was no history of hypertension. His general and systemic examination was normal. Neurological examination was normal, except bilateral 6<sup>th</sup> nerve paresis. Magnetic resonance imaging (MRI) of the brain showed a large well-defined hyperdense contrast-enhancing lesion involving the clivus. In view of the enlarging size of the

Access this article online	
Quick Response Code:	Website: www.asianjns.org
	DOI: 10.4103/1793-5482.131078

#### **Address for correspondence:**

Dr. Amit Agrawal, Department of Neurosurgery, Narayana Medical College Hospital, Chinthareddypalem, Nellore - 524 003, Andhra Pradesh, India. E-mail: dramitagrawal@gmail.com

Discussion

in some foci [Figures 2 and 3].

Primary GCTs of the clivus are a rare lesion with only few reported cases in the literature.<sup>[4-6]</sup> The clinical features of these tumors depend on the location of cranial lesion and symptoms vary in accordance with their actual location. GCT of the sphenoid can present with headache, visual field defects, blindness, diplopia, second through 8th cranial nerve dysfunction, endocrinopathy, and altered mental status;[4-6] on the contrary, temporal bone tumors can present with pain behind the ear, deafness, and facial weakness.<sup>[7]</sup> GCT is characterized by vascularized tissue that contains numerous cytologically benign multinucleated giant cells dispersed through plump, spindly, and/or ovoid cells.<sup>[1,8]</sup> Nuclei of the cells are generally hypochromatic with inconspicuous nucleoli and uncommon mitotic figures.<sup>[9]</sup> Presence of epithelioid histiocytes is rare in GCT and it was described in case of GCT of the tendon sheath, which was largely composed of epithelioid; histiocytes are very rare and in the reported case, the tumor was composed of cellular and hypocellular zones, celluar

lesion and increase in headache, the patient was planned for resurgery [Figure 1]. The patient underwent extended bifrontal

craniontomy and orbitoomy, subfrontal approach, and

decompression of the tumor. Postoperatively was uneventful,

the headache improved but diplopia and 6th nerve paresis

was persisting. Microscopically, the lesion is composed of

multinucleated giant cells admixed with mononuclear stromal

cells. The stromal cells are polygonal, and some of them are elongated spindle-shaped. The cells have vesicular round to

oval nuclei with prominent nucleoli in some of them. Increased vacularity was noted in some areas of tumor. Few foci showed

sheets of clear histiocytes having small round nuclei and clear

cytoplasm. Bony trabeculae rimmed by osteoblasts are noted

#### Agrawal, et al.: Giant cell tumor



Figure 1: MRI of the brain showing extensive lesion involving the clivus



**Figure 2:** (a) Sheets of histiocytes with small round nuclei and clear cytoplasm (H and E, ×100) and (b) Sheets of histiocytes with small round nuclei and clear cytoplasm (H and E, ×400)



Figure 3: Tumor showing multiple osteoclast-like giant cells admixed with stromal cells (H and E,  $\times$ 100)

zone was composed of spindle cells and osteoclast-like giant cells, and the hypocelluar zone was composed of epithelioid clear histiocytes, and it was suggested that the epithelioid histiocytes were the neoplastic cells.<sup>[10]</sup> GCTs needs to be

differentiated from other lesion including chondrosarcoma and chordomas, aneurysmal bone cyst, giant cell reparative granuloma, "Brown tumor" of hyperparathyroidism, and fibrous dysplasia.<sup>[9]</sup> Skull X-rays and angiography have been the traditional investigations for the diagnosis of the GCTs of the skull.<sup>[3,9]</sup> Recently, computed tomography (CT) and MRI have been increasingly used for the diagnosis of these lesions, and the CT appearance of GCT is that of a homogeneous hyperdense mass highly enhancing after contrast administration.<sup>[2,11]</sup> Bony erosions can also be demonstrated by CT scan examination,<sup>[3,11]</sup> and the bone adjacent the lesion can appear hyperplastic in some cases.<sup>[12]</sup> The treatment of choice of GCTs is complete surgical excision and if it can be achieved it is curative; however, as was seen in the present case, it is may not always be feasible.<sup>[9,11,12]</sup> Although controversial, in cases of unresectable tumors or with incomplete excision, radiotherapy remains the other option.<sup>[2,4,7,9,11]</sup>

#### **References**

- Dahlin DC. Caldwell lecture. Giant cell tumor of bone: Highlights of 407 cases. AJR Am J Roentgenol 1985;144:955-60.
- Pai SB, Lalitha RM, Prasad K, Rao SG, Harish K. Giant cell tumor of the temporal bone: A case report. BMC Ear Nose Throat Disord 2005;5:8.
- Vernet O, Ducrey N, Deruaz JP, de Tribolet N. Giant cell tumor of the orbit. Neurosurgery 1993;32:848-51.
- 4. Gupta R, Mohindra S, Mahore A, Mathuriya SN, Radotra BD. Giant cell tumour of the clivus. Br J Neurosurg 2008;22:447-9.
- Zhang QH, Kong F, Yan B, Ni ZL, Lu HL. Endoscopic endonasal surgery for clival tumor. Zhonghua Er Bi Yan Hou Tou Jing Wai Ke Za Zhi 2007;42:7-10.
- Zorlu F, Selek U, Soylemezoglu F, Oge K. Malignant giant cell tumor of the skull base originating from clivus and sphenoid bone. J Neurooncol 2006;76:149-52.
- 7. Pradhan S, Datta NR, Krishnani N, Ayyagari S, Tandon P. Giant cell tumour of the petrous bone. Indian J Cancer 1991;28:177-80.
- Silvers AR, Som PM, Brandwein M, Chong JL, Shah D. The role of imaging in the diagnosis of giant cell tumor of the skull base. AJNR Am J Neuroradiol 1996;17:1392-5.
- Wolfe JT 3<sup>rd</sup>, Scheithauer BW, Dahlin DC. Giant-cell tumor of the sphenoid bone. Review of 10 cases. J Neurosurg 1983;59:322-7.
- Terada T. Giant cell tumor of the tendon sheath composed largely of epithelioid histiocytes. Int J Clin Exp Pathol 2012;5:374-6.
- Findlay JM, Chiasson D, Hudson AR, Chui M. Giant-cell tumor of the middle cranial fossa. Case report. J Neurosurg 1987;66:924-8.
- Watkins LD, Uttley D, Archer DJ, Wilkins P, Plowman N. Giant cell tumors of the sphenoid bone. Neurosurgery 1992;30:576-81.

How to cite this article: Agrawal A, Gali R, Shanthi V, Ramakrishna BA, Mohan KVM. Giant cell tumor of the clivus with presence of epithelioid histiocytes. Asian J Neurosurg 2014;9:48-9. Source of Support: Nil, Conflict of Interest: None declared.

49