

## Mucosal gastric metastases: A very rare site of metastasis from germ cell tumour

DOI: 10.4103/2278-330X.173172

Dear Editor,

Secondary gastric cancer is a rare event identified in <2% of cancer patients. However it may be involved by hematogenous spread from a distant primary (most commonly breast, melanoma and lung) apart from contiguous spread from an adjacent malignancy. Metastasis from testicular germ cell tumours (GCTs) to the stomach is even rare.

A 29-year-old male patient presented with history of orchiectomy in July, 2012 for undescended left testicle and histopathology was suggestive of nonseminomatous GCT (NSGCT) (embryonal with yolk sac element), for which he took no treatment. Computed tomography (CT) chest and abdomen done after 3 months showed liver and lung lesions along with multiple left paraaortic lymphnodes. Serum beta-human chorionic gonadotropin (HCG) was 27650 mIU/ml and serum alpha fetoprotein (AFP) was 115 ng/ml. After 4 cycles of bleomycin, etoposide and cisplatin based chemotherapy till February 2013, repeat scanning showed partial response with normal tumour markers. After 4 months patient again presented with progressive lesions on positron emission tomography-CT with very high level of beta-HCG (24000 mIU/ml) and AFP level of 2.79. Patient was started on paclitaxel, ifosfamide and cisplatin (TIP) based chemotherapy but after 3 cycles of chemotherapy he developed brain metastasis, progressive lung and liver lesions with elevated tumour markers (Beta-HCG 2, 26, 220 mIU/ml and AFP 1.88 ng/ml). Whole brain RT was given for brain metastases. After 2 weeks he presented with hematemesis. Upper gastrointestinal (GI) endoscopy showed multiple gastric lesions suggestive of metastasis. Gastric biopsy was suggestive of metastasis from NSGCT [Figure 1]. In view of progressive disease he was started on gemcitabine and oxaliplatin based chemotherapy.

metastases. After 2 weeks he presented with hematemesis. Upper gastrointestinal (GI) endoscopy showed multiple gastric lesions suggestive of metastasis. Gastric biopsy was suggestive of metastasis from NSGCT [Figure 1]. In view of progressive disease he was started on gemcitabine and oxaliplatin based chemotherapy.

## Discussion

The involvement of the stomach by metastases is rare with the most common reported primaries include breast and lung cancer. The estimated incidence of gastric metastases at autopsy in individuals with a known malignancy varies from 1.7% to 5.4%.<sup>[1]</sup> Most of the individuals harboring such metastases are symptomatic, most commonly with bleeding, pain, vomiting and anorexia.<sup>[2]</sup>

It is estimated that half of the patients with testicular GCTs will have metastases at diagnosis; most common destinations being brain, lymph nodes, liver, and lung.<sup>[3]</sup> Metastases are relatively uncommon to the stomach, adrenals and spleen.




Figure 1: Microphotograph showing submucosa of gastric biopsy infiltrated by a metastatic germ cell tumor (arrow) with areas of haemorrhage (H and E, x100)

In a case series report by Campoli *et al.*, among 771 patients with gastric neoplasms only 20 cases of gastric metastasis were reported with only one case of testicular GCT (Embryonal carcinoma).<sup>[4]</sup> In a retrospective review of five cases with advanced germ cell malignancy (testicular two and retroperitoneal three) and secondaries involving GI tract, 4 patients had duodenal and 1 patient had esophageal involvement. Hemorrhage due to such metastasis could be massive and even fatal.<sup>[5]</sup>

Gastric metastases may be recognizable as abnormalities on gastroscopy (single lesion in the gastric body or by multiple lesions. Often the lesion is described as a “volcano-like” ulcer.<sup>[1,2]</sup> In this case there were multiple well demarcated lesions typical of secondaries seen on endoscopy.

The outcome of the patients of metastatic GCT with GI involvement has been shown to be quite poor, and in one series, 4/5 patients died from unresponsive or progressive disease. Management of such patients mostly requires multimodality approach, including endoscopic techniques, chemotherapy and angio-embolisation.<sup>[5]</sup>

#### Financial support and sponsorship

Nil.

#### Conflicts of interest

There are no conflicts of interest.

**Pankaj Goyal, Amitabh Kumar Upadhyay, D. Chandragouda, J. Gandhi, M. Agarwal**

Departments of Medical Oncology and <sup>1</sup>Pathology, Rajiv Gandhi Cancer Institute and Research Centre, Rohini, New Delhi, India  
**Correspondence to:** Dr. Pankaj Goyal,  
 E-mail: pankaj155@yahoo.com

#### References

1. Oda PG, Kondo H, Yamao T, Saito D, Ono H, Gotoda T, *et al.* Metastatic tumors to the stomach: Analysis of 54 patients diagnosed at endoscopy and 347 autopsy cases. *Endoscopy* 2001;33:507-10.
2. Hamilton SR, Aaltonen LA, editors. World Health Organization Classification of Tumours. Pathology and Genetics of Tumours of the Digestive System. Lyon: IARC Press; 2000.
3. Harikumar R, Harish K, Aravindan KP, Thomas V. Testicular choriocarcinoma with gastric metastasis presenting as hematemesis. *Indian J Gastroenterol* 2004;23:223-4.
4. Campoli PM, Ejima FH, Cardoso DM, Silva OO, Santana Filho JB, Queiroz Barreto PA, *et al.* Metastatic cancer to the stomach. *Gastric Cancer* 2006;9:19-25.
5. Nord C, Fosså SD, Giercksky KE. Gastrointestinal presentation of germ cell malignancy. *Eur Urol* 2000;38:721-4.

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

**How to cite this article:** Goyal P, Upadhyay AK, Chandragouda D, Gandhi J, Agarwal M. Mucosal gastric metastases: A very rare site of metastasis from germ cell tumour. *South Asian J Cancer* 2015;4:153-4.