## **Original Article**

# Transpalpebral Approach for Microsurgical Removal of Tuberculum Sellae Meningiomas

#### Abstract

Background: The evolution of skull base approaches associated with individualization of surgical corridor and minimizing the collateral damage. Achieving the radical removal of tumor and preserving the neurological status of the patient is possible, both with the traditional approaches and keyhole approaches. Our work presents experience using the transpalpebral approach (TPA) for microsurgical removal of tuberculum sellae meningioma (TSM). Materials and Methods: A total of 15 patients with meningiomas underwent microsurgical removal of TSM through TPA. Ten patients were women and five were men. The standard preoperative diagnostic protocol includes magnetic resonance imaging with contrast enhancement, brain computed tomography for neuronavigation. We assess surgical complications, functional and cosmetic outcomes, and surgical parameters, including the time of surgery and intraoperative blood loss. Results: Visual impairment was finding in 100% patients, including slight decrease of vision (46,7%, seven patients), partial vision field loss (six patients, 40%), and serious visual impairment (two patients 13.3%). Visual improvement was noted in ten cases (66.7%), there was no improvement in four cases (26.7%), and one case (6.6%) had transient visual worsening for 4 days and slow improvement in 1 month. Headache disappeared in three patients (50%). There were no cases of cerebrospinal fluid leak. Transient frontal hypoesthesia was noted in all patients (100%) without permanent deficit. Transient palsy of the frontal muscle was noted in four patients for 4-6 months. Histological examination revealed WHO Grade I meningioma in 14 cases and in 1 case WHO Grade II meningioma. No deaths were identified in follow-up at 12 months. The average value of the Modified Rankin Scale was 1.4. The mean length of stay in hospital was 5. Conclusion: TPA is technically difficult and requires some experience to work in deep structures in a small surgical corridor. This technique can be good alternative to traditional fronto-lateral, supraorbital keyhole craniotomies, and endoscopic endonasal approaches.

**Keywords:** Keyhole surgery, minimally invasive neurosurgery, transpalpebral approach, tuberculum sellae meningioma

## Introduction

Tuberculum sellae meningiomas (TSMs) account for about 5%–10% of all intracranial meningiomas.<sup>[1]</sup>

It is well known that the surgical approach to the most TSMs was performed through a large visible frontotemporal skin incision with extended temporal muscle dissection and frontotemporal craniotomy with or without supraorbital bar removal. The evolution of skull base approaches is associated with individualization of surgical corridor and minimizing the collateral damage. For several decades, extended fronto-lateral approaches, such as bifrontal, pterional, and orbitozygomatic were the "working horses" for microsurgical removal of TSM.<sup>[2-8]</sup>

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

Surgical treatment of the TSMs is a complex problem. During this time, it became apparent that the unilateral approach gives the same outcome but with less damage in compared with bifrontal craniotomy.<sup>[9]</sup>

One of the main goals of surgery is the radical removal of the tumor with preserving the visual pathways and choosing the most optimal approach. Achievement of these goals is possible, both with the traditional approaches and keyhole approaches.

The endoscopic removal of TSM is another well-known alternative approach. There are certain disadvantages of the endonasal route: technically more complex, limited angle of view, problems with bleeding control, and a higher risk of postoperative cerebrospinal fluid (CSF) leak.<sup>[10,11]</sup>

How to cite this article: Dzhindzhikhadze RS, Dreval ON, Lazarev VA, Polyakov AV, Kambiev RL, Salyamova EI. Transpalpebral approach for microsurgical removal of tuberculum sellae meningiomas. Asian J Neurosurg 2020;15:98-106. Submission: 22-06-2019 Accepted: 07-10-2019 Published: 25-02-2020 Revaz Semenovich Dzhindzhikhadze<sup>1,2</sup>, Oleg Nikolaevich Dreval<sup>1</sup>, Valeriy Aleksandrovich Lazarev<sup>1</sup>, Andrey Victorovich Polyakov<sup>1,2</sup>, Renat Leonidovich Kambiev<sup>2</sup>, Elvira Igorevna Salyamova<sup>1,2</sup>

<sup>1</sup>Department of Neurosurgery, Russian Medical Academy for Continuing Postgraduate Education, Ministry of Health of Russia, <sup>2</sup>Department of Vascular Neurosurgery, City Clinical Hospital Named After F.I. Inozemtseva, Moscow, Russia

Address for correspondence: Dr. Andrey Victorovich Polyakov, 142115, Moscow Region, Podolsk City, Mashinostroiteley Street, 28-99, Moscow, Russia. E-mail: ap.neurosurg@mail.ru



One of the most important principles of keyhole surgery is not reducing the size of the craniotomy but minimizing the approach-related complications and unnecessary traumatization that is not related to the goal of the surgery. It should be noted that when we use mini approach, we see the same microscopic picture as with extended craniotomies. Adequate patient selection is critical for safe and effective removal of TSM.<sup>[12-15]</sup>

In this report, the authors present results of using the transpalpebral approach (TPA) with miniorbitofrontal craniotomy for microsurgical removal of TSM.

## **Materials and Methods**

A total of 15 patients with meningiomas underwent microsurgical removal of TSM in the Department of Neurosurgery, City Clinical Hospital named after F.I. Inozemtseva, Moscow, Russia, between 2016 and 2018. Ten patients were women, and five were men. All tumors were operated through the transpalpebral incision and keyhole orbitofrontal craniotomy. Summary of 15 cases of TSM is presented in Table 1.

The standard preoperative diagnostic protocol includes magnetic resonance imaging (MRI) with contrast enhancement, brain computed tomography (CT) for neuronavigation, examination by an ophthalmologist and an otolaryngologist, and a routine neurologic examination.

All patients were informed about the alternative extended craniotomies, supraorbital craniotomy with eyebrow skin incision and transsphenoidal endoscopic approach.

We used careful preoperative evaluation of the following parameters: facial and bone anatomy, frontal sinus topography and their relationship with the planning craniotomy, depth of the olfactory groove, and anterior clinoid pneumatization. The potential risk of frontal sinuses opening was evaluated by virtual craniotomy. We also used follow-up assessment of functional and cosmetic outcomes and assessment of disability in accordance with Modified Rankin scale (mRS). Extent of tumor resection was evaluated on a Simpson scale. All parameters were assessed over a period of 12 months.

Surgical parameters, including the time of surgery, intraoperative blood loss, and surgery-related complications, were also reviewed. Postoperatively, all patients underwent MRI with contrast enhancement on 4–5 days and 6–8 months after surgery.

## Surgical technique

Before surgery, the incision is marked with the raised and lowered eyelid for better identification of the natural crease. The patient placed on the operating table with the head fixed in a 3-pin Mayfield head holder and elevated above the heart level, tipping the head downward, and turning in the opposite side from 15° to 30°. Three patients were approached from the left side and 12 from the right side. The side of craniotomy was determined by tumor side lateralization and preoperative dominant side of the visual deficit. The right side is preferable. Before the skin incision, we placed an ophthalmic gel subconjunctivally. Temporary tarsoraphia is performed with a 5-0 nylon suture. The area of the planned incision is infiltrated with an anesthetic solution and vasoconstrictor.

The skin incision is made through the natural fold of the upper eyelid. The incision of the orbicularis oculi muscle is like the projection of the skin incision and subcutaneous tissue [Figure 1]. The dissection of the orbicularis muscle was performed with preservation of the orbital septum and

Table 1: Summary of 15 cases of tuberculum sellae meningiomas											
Case	Gender	Age	Max tumor	Lateralization	Surgical	Blood	Extent of	Recurrence	Visual	Length of	Follow-up
			size (mm)		time (min)	loss (ml)	resection		changes	stay (days)	period (months)
1	Male	54	21.4	Right	200	50	Simpson II	None	Improve	4	10
2	Female	52	32.2	None	240	100	Simpson II	None	Improve	5	17
3	Male	48	20.3	None	175	50	Simpson II	None	Improve	4	8
4	Female	62	30.8	None	210	100	Simpson II	None	None	5	16
5	Female	60	38.1	Right	245	150	Simpson II	None	None	6	21
6	Female	55	25.6	None	215	50	Simpson II	None	Improve	4	18
7	Female	48	30.2	None	245	150	Simpson IV	None	Improve	5	14
8	Male	48	24.2	None	245	50	Simpson II	None	Improve	4	12
9	Male	75	25.1	Right	220	100	Simpson II	None	Improve	5	14
10	Female	69	39.2	Right	275	300	Simpson IV	+	None	7	11
11	Female	60	37.7	Left	230	150	Simpson II	None	None	5	20
12	Female	52	29.9	None	225	150	Simpson II	None	Improve	6	22
13	Female	51	26.1	None	200	50	Simpson II	None	Improve	4	7
14	Male	59	38.7	Left	230	200	Simpson IV	None	Transient	7	24
									worsering		
15	Female	64	30.7	Left	235	100	Simpson II	None	Improve	4	19
· D							1		1		-

+: Rec

lateral canthal ligament. A single musculocutaneous flap is formed. Subperiosteal dissection was performed in the supraorbital region from the lateral margin of the supraorbital notch to the frontal-zygomatic suture. The temporal muscle dissected with monopolar from the place of its attachment at the keyhole-point area and mobilized laterally to visualize the area of burr hole placement. At this step, an important nuance is a careful work with soft tissues to avoid the damage of the facial nerve branches that innervate the orbicularis oculi muscle. According to the work of Ouattara et al. in 63.3% of cases, the muscle is innervated by most of the temporal, upper zygomatic, and lower zygomatic branches and by the upper buccal branches.<sup>[16]</sup> Schmidt et al. reported that lateral border of the orbicularis oculi muscle, where the temporal and zygomatic nerves insert into the muscle, the mean distance between the temporal and zygomatic nerves was  $1.72 \pm 0.62$  cm, and the average horizontal distance from the lateral canthus to the point where the nerve changes from a vertical course to a horizontal course was  $4.70 \pm 0.79$  cm.<sup>[17]</sup>



Figure 1: Projections of skin, subcutaneous tissue, and musculus orbicuaris oculi incision (Black line). Yellow line – projections of facial nerve branches

The one burr hole is placed in the keyhole point with the high-speed diamond drill. After that, a miniorbitofrontal craniotomy is performed, including the roof of the orbit and the portion of the frontal bone about 1-1.5 cm of the frontal process of the zygomatic bone. The diameter of the bone window was not exceeding 2.0-2.5 cm. The roof of the orbit was broken with a chisel [Figure 2a-f].

The dura mater (DM) is opened in a C-shaped fashion. The first important step is early brain relaxation. Basal cisterns are opened with sharp dissection. We use retractorless technique and dynamic retraction to protect the brain and minimizing traction damage. The next step is to identify ipsilateral optic nerve and the internal carotid artery. Small perforators on the inferior surface of the optic nerves and chiasm are preserved. In three cases, the optic canal was unroofed because of its invasion by the tumor. The tumor is devascularized in the early stages. The pituitary stalk is usually displaced posteriorly and easily dissected from the meningioma. In some cases, if the tumor is soft, we use ultrasonic aspirator to debulk the tumor. After removal of the tumor, we use endoscopic assistance to assess the residual volume. The matrix was coagulated, and the basal dura was resected. In eight cases (53.3%), extradural resection of the anterior clinoid process was performed. The DM is tightly closed. The bone flap is fixed with miniplates or CranioFix. Bone cement is optionally used for additional fixation. The wound is sutured layer by layer, an intradermal suture of absorbable 6-0 thread is applied to the skin. In the first 2 h after the operation, cubes of ice are placed on the area of the postoperative wound to reduce the periorbital edema.

#### **Patients population**

Of the 15 patients, there were 10 women (66.7%) and 5 men (33.7%). The female-to-male ratio was 2:1. The mean age was 57.1 years (range: 48–75 years). None of



Figure 2: Steps of performing transpalpebral approach. (a) Skin incision. (b and c) Soft-tissue dissection. (d) Burr hole forming with a high-speed drill. (e and f) Mini-orbitofrontal craniotomy

the patients underwent prior surgery and did not undergo radiation courses.

## **Clinical features**

Visual impairment – the most frequent symptoms that find in 100% patients, including slight decrease of vision (seven patients, 46.7%), partial vision loss (six patients, 40%), and serious visual impairment (two patients, 13.3%).

The other symptoms include headache (six patients, 40%), anosmia (three patients, 20%), and diplopia (one patient, 6.7%). None of the patients had pre- and post-operative endocrinological deficit.

## **Radiological features**

Pre- and post-operative MRI with contrast enhancement was used and combined with CT for craniotomy planning.

The mean tumor diameter and volume were 30.0 mm (range: 20.3-43.2 mm) and  $15.2 \text{ cm}^3$  (range:  $11.3-27.1 \text{ cm}^3$ ). There are four R-lateralization cases and three L-lateralization cases. Tumor calcification was found in two patients (13.3%).

## Results

In three patients, resection had to be subtotal, a residual tumor was observed on follow-up MRI as expected. In three patients, the tumor was attached to the artery of Heubner.

There were no serious approach-related complications (CSF leakage, epidural and subdural hematoma, wound infection, and keloid scar). In four cases, the frontal sinus was opened. In one case, the mucous membrane was damaged, and cranialization was performed with abdominal fat packing with vancomycin and additional fibrin glue sealing. In other three cases, a small frontal sinus defect was waxed.

We used rigid endoscopes (Karl Storz, Germany) with  $0^{\circ}$ - and  $30^{\circ}$ -angled lenses for microsurgery assistance. Simpson II resection achieved in 80% cases. Invasion in the optic canal was in three patients. The mean surgical time was 226 min and the mean blood loss was 110 ml.

All patients were in the intensive care unit for 1 day. An early postoperative CT was performed within 24 h to exclude complications.

Visual improve was noted in ten cases (66.7%), there was no improvement in four cases (26.7%), and one case (6.6%) had transient visual worsening for 4 days and slow improvement in 1 month. Our results in part of the visual deficit are comparable with the results of other authors. Obviously, the degree of changes in visual impairments depends on many factors that not directly related to the surgical approach (duration of the symptoms, tumor size and its adherence to the optic nerves, and tumor location).

There were no motor or sensory limb dysfunctions in the postoperative period. All parameters are shown in Table 2. Headache disappeared in three patients (50%). There were no cases of CSF leak.

Two major cosmetic outcomes of TPA were also evaluated. Transient frontal hypoesthesia for 3–5 months was reported in all patients (100%), permanent deficit was not reported. Transient palsy of the frontal muscle for 6–8 months was reported for four patients, permanent palsy was not reported.

Histological examination revealed WHO Grade I meningioma in 14 cases and in 1 case WHO Grade II meningioma. Almost 80% of patients have Simpson II resection. No deaths occurred in follow-up at 12 months. The average value of mRS was 1.4. The mean length of stay in hospital was 5. The mean follow-up period was 15.5 months. No patients had endocrinological disturbances. The list of complications is presented in Table 3. Satisfaction with the postoperative cosmetic outcome after TPA is shown in Table 4.

Clinical case of using TPA for microsurgical removing of TSM is shown in Figures 3 and 4.

Cosmetic outcomes are shown in Figure 5. The patients have consented to the submission of this article for submission to the journal.

## Discussion

One of the first article, devoted to the description of suprasellar meningiomas, was the report of Cushing and Eisenhardt in 1929.<sup>[3]</sup> Various approaches were used in the surgical treatment of this pathology, using the bifrontal and unilateral approaches with different series of patients.<sup>[18-23]</sup>

Despite the advantages of traditional approaches, there still exist some risks of approach-related complications that is not related with the surgical target: atrophy of the temporal muscle, asymmetry of the face, risk of dysfunction of the temporomandibular joint, pain and numbness in the orofacial region, pain during chewing, discomfort in wearing glasses, injury of the frontal branch of the facial nerve, alopecia in the postoperative scar, risk of epidural hematoma, and retraction-induced damage.<sup>[24-29]</sup>

Over time, it became obvious, that the microsurgical removal of TSM possible through keyhole approaches with minimum risks of approach-related complications. Minimally invasive neurosurgery allows us to adapt and individualize the surgical view and reduces the associated surgical trauma. The concept of keyhole surgery was popularized by Perneczky and Reisch. Many authors presented effective and safety outcomes of mini approaches (minisupraorbital craniotomy) for the parasellar and anterior cranial fossa lesions with adequate patient selection.<sup>[12,13,30-37]</sup>



Figure 3: Preoperative planning transpalpebral approach. (a-c). Preoperative magnetic resonance imaging with cerebrospinal fluid, tuberculum sellae meningioma. (d) Projection line of skin incision (arrow). (e and f) Virtual craniotomy, planning of transpalpebral approach in computer model. (g and h) Using the neuronavigation to mark the frontal sinus



Figure 4: Intraoperative pictures, microsurgical removal of tuberculum sellae meningioma through transpalpebral approach. (a) Dural opening. (b) Basal cistern opening, brain relaxation. (c) Tumor debulking with ultrasonic aspirator. (d) Tumor resecting partially with microscissors. (e) Part of meningioma removing through small craniotomy. (f) Endoscope assistance. (g-i) 1-month postoperative magnetic resonance imaging with cerebrospinal fluid

We have an experience of using TPA in the skull base surgery. Our cases include performing TPA for anterior circulation aneurysms, microsurgical removal tumors of the anterior cranial fossa, and orbital cavernomas.



Figure 5: (a-d) Cosmetic outcomes after transpalpebral approach

Table 3: Postoperative complications aftertranspalpebral approach						
	complications, n (%)					
Wound infection	0					
CSF leakage	0					
Epidural, subdural hematoma	0					
Traction damage	2 (13.3)					
Endocrinologic complications	0					
Postoperative seizure	0					
Cognitive worsening	3 (20)					
Keloid scar	0					
Lethal cases in 12 months follow-up	0					
CSE Comphragning I fluid						

CSF – Cerebrospinal fluid

 Table 4: Satisfaction with the postoperative cosmetic outcome after transpalpebral approach in 15 patients

 Scale of satisfaction

 Number of cases, n (%)

Scale of satisfaction	Number of cases, n (70
Unsatisfactory	0
Good	2 (13.3)
Excellent	13 (86.7)

We assessed the functional and cosmetic outcomes, surgical complications and concluded that TPA could be successfully performed as an alternative method to traditional extended approaches and endoscopic routes in the skull base surgery.<sup>[38-40]</sup>

In this report, we presented outcomes in 15 patients diagnosed with TSM. A surgical approach was selected with critical assessment of facial and bone anatomy, sizes of the tumor.

Visual disturbances are the main symptom of TSMs. It is agreed that the visual outcome depends on the duration of the disease and the visual deficit before surgery.<sup>[41]</sup> One of the most important goals is to decrease additional damage to the already impaired visual pathways. Some reports showing improvement of visual function in a wide range from 25% to 80%.<sup>[42-49]</sup> Mathiesen and Kihlström hypothesizing that early optic canal decompression and optic nerve release can improve visual outcomes.<sup>[50]</sup>

Al-Mefty and Smith report that direct compression of the tumor on the vascular support to the optic chiasm and optic nerve is the main cause of visual impairment.<sup>[43]</sup> Some studies compared the endoscopic transsphenoidal versus supraorbital keyhole approach for the resection of TSM.<sup>[51,52]</sup> However, adequate minimization of approach allows us to achieve similar results on the visual pathways, without increasing the risk of developing approach-related complications.

It is known that the mortality rate for microsurgical removing of TSM varies from 0 to 8.7%.<sup>[41,45,48,49]</sup> According to the report by Nakamura *et al.*, there is a definite dependence of mortality on the approach performed (9.5% bifrontal craniotomy vs. 0% pterional/fronto-lateral craniotomy).<sup>[9]</sup> The outcome of the TSM surgery is dependent more on the clinical state of the patient. Individualization and minimization of the approach allow as to reduce the damage of normal structures that are not related to the pathology.

Subtotal resection has ranged from 8% to 33%.<sup>[9,22,41,49]</sup> The degree of tumor resection depends primarily on the experience of the neurosurgeon. In the report of Nakamura *et al.*, there were no significant differences in the degree of tumor resection and the chosen approach.<sup>[9]</sup> It is always necessary to assess the relationship of the tumor with the vessels and optic nerves. If necessary, the residual part is left in critical structures to avoid the neurological deficit and it is not associated with transcranial approach (extended or keyhole craniotomy).

CSF leakage complications range from 4% to 33%.<sup>[9,23,45,53,54]</sup> Linsler *et al.* reported that CSF leakage was manifested after supraorbital keyhole approach (6%), in another article by Fatemi *et al.* CSF leak occurred after transnasal endoscopic removal of the tumor (29%).<sup>[51,52]</sup>

There are few reports of the development of endocrinological complications-diabetes insipidus with a frequency of 5%-33%.<sup>[19,23,43,47]</sup> There are some reports on the development of endocrinological complications in single cases, both after endonasal approach and supraorbital keyhole craniotomy.<sup>[51,52]</sup>

TPA planning should be carried out in advance with using virtual craniotomy. When we choose TPA, it is important to assess the individual surgical route to the lesion and adequately minimize craniotomy. This approach gives a surgical view such as pterional, lateral supraorbital, and orbitozygomatic craniotomy. As we have a predefined subfrontal route to the lesion, it is not necessary for the large exposure of the cerebral cortex.[55] Some modifications of the TPA: resection of the orbital roof, extradural resection of the anterior clinoid, and partially sphenoidal wing resection could be successfully applied to increase the microsurgical corridors. Creating a comfortable work through small craniotomy by opening basal cisterns can significantly increase the space for manipulating with microsurgical instruments and removing large volume tumors. Using of neuronavigation helps to determine the lateral border of the frontal sinus. We individualize and determine the surgical corridor because any damaging of the frontal sinus increases the risk of infectious complications and CSF leakage.<sup>[56,57]</sup>

## Conclusion

We conclude that the TPA can be used for removing TSM with good cosmetic and functional outcomes.

Over the entire microsurgical era, many approaches have been used to the TSM. Over time, they underwent a certain evolution from extended approaches to mini-approaches and endonasal routes. Each craniotomy has specific advantages and disadvantages. It is important that TPA is technically more difficult and requires some experience to work in a deep structure through a small surgical corridor. This technique can be good alternative to traditional fronto-lateral craniotomies, eyebrow supraorbital and endonasal approaches.

## **Declaration of patient consent**

The authors certify that they have obtained all appropriate patient consent forms. In the form, the patient has given their consent for their images and other clinical information to be reported in the journal. The patient understands that name and initials will not be published and due efforts will be made to conceal identity, but anonymity cannot be guaranteed.

## Financial support and sponsorship

Nil.

#### **Conflicts of interest**

There are no conflicts of interest.

#### References

- Ruggeri AG, Cappelletti M, Fazzolari B, Marotta N, Delfini R. Frontobasal midline meningiomas: Is it right to shed doubt on the transcranial approaches? Updates and review of the literature. World Neurosurg 2016;88:374-82.
- 2. Cushing H. The Pituitary Body and its Disorders: Clinical States Reduced by Disorders of the Hypophysis Cerebri. Philadelphia:

JB Lippincott; 1912.

- Cushing H, Eisenhardt L. Meningiomas Arising from the Tuberculum Sellae: With the Syndrome of Primary Optic Atrophy and Bitemporal Field Defects Combined with a Normal Sellae Turcica in A Middle-Aged Person. Chicago: American Medical Association; 1929.
- Dandy W. A new hypophysis operation. Bull Johns Hopkins Hosp 1918;29:154-5.
- 5. Frazier CH. I. An approach to the hypophysis through the anterior cranial fossa. Ann Surg 1913;57:145-50.
- 6. Heuer G. The surgical approach and the treatment of tumors and other lesions about the optic chiasm. Surg Gynecol Obstet 1931;53:489-518.
- 7. Horsley V. Remarks on the surgery of the central nervous system. Br Med J 1890;2:1286-92.
- 8. Krause F. Hirnchirurgie. Deutsche Lin 1905;8:953-1042.
- Nakamura M, Roser F, Struck M, Vorkapic P, Samii M. Tuberculum sellae meningiomas: Clinical outcome considering different surgical approaches. Neurosurgery 2006;59:1019-28.
- 10. Har-El G, Casiano RR. Endoscopic management of anterior skull base tumors. Otolaryngol Clin North Am 2005;38:133-44, 9.
- Jho HD, Ha HG. Endoscopic endonasal skull base surgery: Part 1 – The midline anterior fossa skull base. Minim Invasive Neurosurg 2004;47:1-8.
- Perneczky A, Reisch R. Keyhole approaches in neurosurgery. Vol. 1 Concept and Surgical Technique. Vienna: Springer-Verlag Wien; 2008.
- Reisch R, Perneczky A. Ten-year experience with the supraorbital subfrontal approach through an eyebrow skin incision. Neurosurgery 2005;57:242-55.
- 14. Teo C, Sugrhue M. Principles and practice of keyhole brain surgery. Stuttgart: Georg Thieme Verlag; 2015.
- 15. van Lindert E, Perneczky A, Fries G, Pierangeli E. The supraorbital keyhole approach to supratentorial aneurysms: Concept and technique. Surg Neurol 1998;49:481-9.
- Ouattara D, Vacher C, de Vasconcellos JJ, Kassanyou S, Gnanazan G, N'Guessan B. Anatomical study of the variations in innervation of the orbicularis oculi by the facial nerve. Surg Radiol Anat 2004;26:51-3.
- 17. Schmidt BL, Pogrel MA, Hakim-Faal Z. The course of the temporal branch of the facial nerve in the periorbital region. J Oral Maxillofac Surg 2001;59:178-84.
- Brihaye J, Brihaye-van Geertruyden M. Management and surgical outcome of suprasellar meningiomas. Acta Neurochir Suppl (Wien) 1988;42:124-9.
- Ehlers N, Malmros R. The suprasellar meningioma. A review of the literature and presentation of a series of 31 cases. Acta Ophthalmol Suppl 1973;121:1-74.
- 20. Gökalp HZ, Arasil E, Kanpolat Y, Balim T. Meningiomas of the tuberculum sella. Neurosurg Rev 1993;16:111-4.
- Olivecrona H, Tönnis W. The suprasellar meningiomas. Handbuch der Neurochirurgie. Berlin: Springer-Verlag; 1967. p. 167-72.
- 22. Solero CL, Giombini S, Morello G. Suprasellar and olfactory meningiomas. Report on a series of 153 personal cases. Acta Neurochir (Wien) 1983;67:181-94.
- 23. Symon L, Rosenstein J. Surgical management of suprasellar meningioma. Part 1: The influence of tumor size, duration of symptoms, and microsurgery on surgical outcome in 101 consecutive cases. J Neurosurg 1984;61:633-41.
- Boari N, Spina A, Giudice L, Gorgoni F, Bailo M, Mortini P. Fronto-orbitozygomatic approach: Functional and cosmetic outcomes in a series of 169 patients. J Neurosurg

2018;128:466-74.

- 25. Brazoloto TM, de Siqueira SR, Rocha-Filho PA, Figueiredo EG, Teixeira MJ, de Siqueira JT, *et al.* Post-operative orofacial pain, temporomandibular dysfunction and trigeminal sensitivity after recent pterional craniotomy: Preliminary study. Acta Neurochir (Wien) 2017;159:799-805.
- 26. Chalouhi N, Jabbour P, Ibrahim I, Starke RM, Younes P, El Hage G, *et al.* Surgical treatment of ruptured anterior circulation aneurysms: Comparison of pterional and supraorbital keyhole approaches. Neurosurgery 2013;72:437-41.
- 27. de Andrade Júnior FC, de Andrade FC, de Araujo Filho CM, Carcagnolo Filho J. Dysfunction of the temporalis muscle after pterional craniotomy for intracranial aneurysms. Comparative, prospective and randomized study of one flap versus two flaps dieresis. Arq Neuropsiquiatr 1998;56:200-5.
- Soleman J, Leiggener C, Schlaeppi AJ, Kienzler J, Fathi AR, Fandino J. The extended subfrontal and fronto-orbito-zygomatic approach in skull base meningioma surgery: Clinical, radiologic, and cosmetic outcome. J Craniofac Surg 2016;27:433-40.
- Welling LC, Figueiredo EG, Wen HT, Gomes MQ, Bor-Seng-Shu E, Casarolli C, *et al.* Prospective randomized study comparing clinical, functional, and aesthetic results of minipterional and classic pterional craniotomies. J Neurosurg 2015;122:1012-9.
- Chen YH, Lin SZ, Chiang YH, Ju DT, Liu MY, Chen GJ. Supraorbital keyhole surgery for optic nerve decompression and dura repair. J Neurotrauma 2004;21:976-81.
- Ditzel Filho LF, McLaughlin N, Bresson D, Solari D, Kassam AB, Kelly DF. Supraorbital eyebrow craniotomy for removal of intraaxial frontal brain tumors: A technical note. World Neurosurg 2014;81:348-56.
- 32. Dlouhy BJ, Chae MP, Teo C. The supraorbital eyebrow approach in children: Clinical outcomes, cosmetic results, and complications. J Neurosurg Pediatr 2015;15:12-9.
- Fischer G, Stadie A, Reisch R, Hopf NJ, Fries G, Böcher-Schwarz H, *et al.* The keyhole concept in aneurysm surgery: Results of the past 20 years. Neurosurgery 2011;68:45-51.
- Ivan ME, Lawton MT. Mini supraorbital approach to inferior frontal lobe cavernous malformations: Case series. J Neurol Surg A Cent Eur Neurosurg 2013;74:187-91.
- 35. Reisch R, Perneczky A, Filippi R. Surgical technique of the supraorbital key-hole craniotomy. Surg Neurol 2003;59:223-7.
- Szabo KA, Cheshier SH, Kalani MY, Kim JW, Guzman R. Supraorbital approach for repair of open anterior skull base fracture. J Neurosurg Pediatr 2008;2:420-3.
- Wilson DA, Duong H, Teo C, Kelly DF. The supraorbital endoscopic approach for tumors. World Neurosurg 2014;82:e243-56.
- Dzhindzhikhadze RS, Dreval' ON, Lazarev VA, Polyakov AV. The transpalpebral keyhole approach in surgery of orbital cavernomas: A case report and literature review. Zh Vopr Neirokhir Im N N Burdenko 2018;82:73-80.
- Dzhindzhikhadze RS, Dreval ON, Lazarev VA, Polyakov AV. Transpalpebral approach in skull base surgery: How I do it. Acta Neurochir (Wien) 2019;161:133-7.
- 40. Dzhindzhikhadze RS, Dreval' ON, Lazarev VA, Polyakov AV,

Kambiev RL. Transpalpebral craniotomy in skull base surgery. Zh Vopr Neirokhir Im N N Burdenko 2018;82:48-58.

- Goel A, Muzumdar D, Desai KI. Tuberculum sellae meningioma: A report on management on the basis of a surgical experience with 70 patients. Neurosurgery 2002;51:1358-63.
- 42. Al-Mefty O, Holoubi A, Rifai A, Fox JL. Microsurgical removal of suprasellar meningiomas. Neurosurgery 1985;16:364-72.
- Al-Mefty O, Smith RR. Tuberculum sellae meningiomas. In: Al-Mefty O, editor. Meningiomas. New York: Raven Press, Ltd.; 1991. p. 395-411.
- 44. Andrews BT, Wilson CB. Suprasellar meningiomas: The effect of tumor location on postoperative visual outcome. J Neurosurg 1988;69:523-8.
- 45. Arai H, Sato K, Okuda O, Miyajima M, Hishii M, Nakanishi H, *et al.* Transcranial transsphenoidal approach for tuberculum sellae meningiomas. Acta Neurochir (Wien) 2000;142:751-6.
- Ciric I, Rosenblatt S. Suprasellar meningiomas. Neurosurgery 2001;49:1372-7.
- Conforti P, Moraci A, Albanese V, Rotondo M, Parlato C. Microsurgical management of suprasellar and intraventricular meningiomas. Neurochirurgia (Stuttg) 1991;34:85-9.
- Fahlbusch R, Schott W. Pterional surgery of meningiomas of the tuberculum sellae and planum sphenoidale: Surgical results with special consideration of ophthalmological and endocrinological outcomes. J Neurosurg 2002;96:235-43.
- Jallo GI, Benjamin V. Tuberculum sellae meningiomas: Microsurgical anatomy and surgical technique. Neurosurgery 2002;51:1432-39.
- Mathiesen T, Kihlström L. Visual outcome of tuberculum sellae meningiomas after extradural optic nerve decompression. Neurosurgery 2006;59:570-6.
- Fatemi N, Dusick JR, de Paiva Neto MA, Malkasian D, Kelly DF. Endonasal versus supraorbital keyhole removal of craniopharyngiomas and tuberculum sellae meningiomas. Neurosurgery 2009;64:269-84.
- 52. Linsler S, Fischer G, Skliarenko V, Stadie A, Oertel J. Endoscopic assisted supraorbital keyhole approach or endoscopic endonasal approach in cases of tuberculum sellae meningioma: Which surgical route should be favored? World Neurosurg 2017;104:601-11.
- Galal A, Faisal A, Al-Werdany M, El Shehaby A, Lotfy T, Moharram H, *et al.* Determinants of postoperative visual recovery in suprasellar meningiomas. Acta Neurochir (Wien) 2010;152:69-77.
- Terasaka S, Asaoka K, Kobayashi H, Yamaguchi S. Anterior interhemispheric approach for tuberculum sellae meningioma. Neurosurgery 2011;68:84-8.
- Owusu Boahene KD, Lim M, Chu E, Quinones-Hinojosa A. Transpalpebral orbitofrontal craniotomy: A minimally invasive approach to anterior cranial vault lesions. Skull Base 2010;20:237-44.
- 56. Andersen NB, Bovim G, Sjaastad O. The frontotemporal peripheral nerves. Topographic variations of the supraorbital, supratrochlear and auriculotemporal nerves and their possible clinical significance. Surg Radiol Anat 2001;23:97-104.
- Kazkayasi M, Batay F, Bademci G, Bengi O, Tekdemir I. The morphometric and cephalometric study of anterior cranial landmarks for surgery. Minim Invasive Neurosurg 2008;51:21-5.