

## Case Report

## Glioma Simultaneously Present with Adjacent Meningioma: Case Report and Literature Review

## Abstract

A 51-year-old male patient presented to us with an episode of generalized tonic-clonic seizure. Magnetic resonance imaging revealed a dural-based contrast-enhancing lesion in the right temporal lobe and another heterogeneously contrast-enhancing intra-axial lesion in the right insula adjacent to it. Histopathology confirmed it as a meningioma adjacent to an anaplastic oligodendroglioma. This is only the second such case reported in literature. Literature on “adjacent site” gliomas and meningioma was also reviewed.

**Keywords:** Double tumors, glioma, meningioma

**Satyashiva Munjal,  
Jitendra Kumar,  
Sunila Jain<sup>1</sup>,  
Veer Singh Mehta**

*Department of Neurosurgery, Paras Hospital, Gurgaon, Haryana, <sup>1</sup>Department of Histopathology, Sir Ganga Ram Hospital, New Delhi, India*

## Introduction

Multiple primary brain tumors are usually seen in the presence of genetic disorders such as neurofibromatosis, tuberous sclerosis, or in patients who have received radiotherapy.<sup>[1]</sup> In the absence of such factors, multiple primary brain tumors, especially at adjacent sites, are rare. We present a case of a 51-year-old male patient who presented to us after a single episode of generalized tonic-clonic seizure and was finally diagnosed with “adjacent site” anaplastic oligodendroglioma and meningioma. This is only the second such case reported in literature.

## Case Report

A 51-year-old male presented to us with a complaint of one episode of generalized tonic-clonic seizure 15 days ago. The seizure was associated with loss of consciousness for 10 min and jerky movements of all four limbs. There was no history of aura, postictal headache or weakness, tongue bite, or loss of continence associated with it. On examination, he had no neurological deficit. There were no neurocutaneous markers. Magnetic resonance imaging brain [Figure 1] revealed an ill-defined area of altered signal intensity in the right frontotemporal and insular region which was hypointense on T1, hyperintense on T2 with patchy areas

of enhancement suggestive of a high-grade glioma. Another avidly enhancing lesion dural-based extra-axial lesion was overlying right temporal lobe (adjacent to the other lesion) suggestive of a meningioma.

Under neuronavigation, the patient underwent the right frontotemporal craniotomy. After reflecting the dural flap anteroinferiorly, a dural-based lesion was seen attached to the temporal dura and extending into sylvian fissure suggestive of meningioma [Figure 2]. The sylvian fissure was opened up due to lesion. The other tumor was intra-axial, soft, grey-white, suckable, moderately vascular, and extended into frontal operculum and temporal lobe around the right middle cerebral artery suggestive of high-grade glioma. Postoperative period was uneventful, and the patient was discharged on postoperative day 6. On histopathological examination [Figure 3], the dural-based lesion revealed a tumor composed of sheets and whorls of spindle cells. There was mild nuclear atypia. Few psammoma bodies were also present. These findings were suggestive of Grade 1 meningioma. The intra-axial lesion revealed a moderately cellular tumor composed of sheets of tumor cells with a fibrillary background. There was moderate nuclear atypia with increased mitotic activity. On immunohistochemistry, tumor cells were positive for glial fibrillary acidic protein, p53. Ki-67 was 12%–15%. The findings were suggestive of anaplastic oligodendroglioma (Grade 3).

## Address for correspondence:

*Dr. Satyashiva Munjal,  
Department of Neurosurgery,  
Paras Hospital,  
Gurgaon - 122 002,  
Haryana, India.  
E-mail: satyashivamunjal@gmail.com*

## Access this article online

**Website:** www.asianjns.org

**DOI:** 10.4103/ajns.AJNS\_308\_17

## Quick Response Code:



**How to cite this article:** Munjal S, Kumar J, Jain S, Mehta VS. Glioma simultaneously present with adjacent meningioma: Case report and literature review. Asian J Neurosurg 2019;14:272-4.

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

**For reprints contact:** reprints@medknow.com


## Discussion

Multiple intracranial tumors are extremely rare. They are more commonly seen in patients with genetic disorders such as neurofibromatosis or tuberous sclerosis.<sup>[1]</sup> They may also be seen in patients who have received radiotherapy treatment.<sup>[1]</sup> The incidence of primary brain tumors with different histology is 0.3% of all brain tumors.<sup>[1]</sup> The most frequently occurring combination is that of meningioma and glioma, followed by meningioma and neurinoma and meningioma and pituitary adenoma.<sup>[1]</sup> Even though meningioma along with glioma is the most frequently occurring combination, its occurrence at adjacent sites has been reported by only


a few authors<sup>[2-10]</sup> [Table 1]. More so, occurrence of an oligodendroglioma adjacent to a meningioma is even rarer, this case being only the second such case available in the English literature.<sup>[2]</sup>

Various hypotheses have been proposed to explain the occurrence of this rare phenomenon. Since meningiomas and gliomas are the most commonly occurring primary brain tumors, most authors feel that this concurrence of meningioma and glioma at adjacent sites is a mere co-incidence.<sup>[1,6,7]</sup> Certain genetic factors, exposure to chemicals, trauma, or some immunological mechanism may be responsible for this phenomenon. Some are of the opinion that a locally acting oncogenic paracrine factor from meningioma may induce malignant transformation in adjacent brain parenchyma and glial cells.<sup>[6]</sup> Similarly, irritative effects from low-grade glioma may induce meningeal proliferation. Exact mechanism is, however, yet to be elucidated.

In such cases, it is usually not possible to make a preoperative diagnosis of “double tumors.” One must be aware of the possibility of such an entity and suspect it if perilesional edema does not correlate with the presence of



**Figure 1:** (a) Magnetic resonance imaging brain T1-weighted image showing a hypointense right insular lesion (asterisk) and an isointense dural-based lesion (arrow). (b and c) Contrast Magnetic resonance imaging image showing strongly enhancing dural-based lesion suggestive of a meningioma (arrow) and an heterogeneously enhancing lesion in right insular region suggestive of a glioma (asterisk)




**Figure 2:** Intraoperative image showing the dural-based meningioma extending into the sylvian fissure (arrow)

**Table 1: Summary of reported cases of “adjacent site” glioma and meningioma**

| Author ( <i>et al.</i> )                    | Year | Age (years) | Sex (male/female) | Location       | Pathology                                    |
|---|------|-------------|-------------------|----------------|--|
| Gass and Van Wagenen <sup>[2]</sup>         | 1950 | 56          | Female            | Right frontal  | ODG + meningioma                             |
| Nagashima <i>et al.</i> <sup>[3]</sup>      | 1963 | 42          | Male              | Left frontal   | Glioma (low grade) + fibroblastic meningioma |
| Strong <i>et al.</i> <sup>[4]</sup>         | 1976 | 56          | Female            | Right frontal  | GBM + meningioma                             |
| Strong <i>et al.</i> <sup>[4]</sup>         | 1976 | 53          | Male              | Lt. frontal    | GBM + syncytial meningioma                   |
| Marra <i>et al.</i> <sup>[5]</sup>          | 1977 | 63          | Male              | Right parietal | GBM + meningotheial meningioma               |
| Goyal <i>et al.</i> <sup>[6]</sup>          | 2003 | 72          | Male              | Right temporal | GBM + fibrous meningioma                     |
| Nestler <i>et al.</i> <sup>[7]</sup>        | 2007 | 49          | Male              | Left frontal   | GBM + fibrous meningioma                     |
| Suzuki <i>et al.</i> <sup>[8]</sup>         | 2010 | 75          | Female            | Left temporal  | GBM + meningotheial meningioma               |
| Chen <i>et al.</i> <sup>[9]</sup>           | 2010 | 63          | Female            | Left frontal   | GBM + fibroblastic meningioma                |
| Shahmohammadi <i>et al.</i> <sup>[10]</sup> | 2016 | 49          | Female            | Right Frontal  | GBM + meningotheial meningioma               |
| Present case                                | 2017 | 51          | Male              | Right temporal | Anaplastic ODG + meningioma (Grade I)        |

ODG – Oligodendroglioma; GBM – Glioblastoma multiforme



**Figure 3:** (a) Highly cellular tumor with perinuclear halos and capillary network in the background. Brisk mitosis (arrow) is seen. (H and E, ×40) suggestive of an anaplastic oligodendroglioma. (b) Ki67 immunohistochemistry showing high proliferative activity. (c) Meningioma with sheets and whorls of tumor cells. Psammoma bodies are seen (H and E, ×20)

a single benign lesion or intraoperatively the tumor does not correlate with preoperative radiology. In certain cases, perilesional edema in meningioma may mask a small low-grade astrocytoma. Biopsy helps to provide tissue diagnosis in such cases.

## Conclusion

Multiple primary brain tumors are rare in the absence of predisposing factors such as genetic disorders or radiotherapy. A surgeon must be aware of the entity “adjacent site” double tumors when radiological and intraoperative findings do not corroborate. Biopsy helps to confirm the diagnosis in these cases.

## Declaration of patient consent

The authors certify that they have obtained all appropriate patient consent forms. In the form the patient(s) has/have given his/her/their consent for his/her/their images and other clinical information to be reported in the journal. The patients understand that their names and initials will not be published

and due efforts will be made to conceal their identity, but anonymity cannot be guaranteed.

## Financial support and sponsorship

Nil.

## Conflicts of interest

There are no conflicts of interest.

## References

1. Spallone A, Santoro A, Palatinsky E, Giunta F. Intracranial meningiomas associated with glial tumours: A review based on 54 selected literature cases from the literature and 3 additional personal cases. *Acta Neurochir (Wien)* 1991;110:133-9.
2. Gass H, Van Wagenen WP. Meningioma and oligodendroglioma adjacent in the brain; case report. *J Neurosurg* 1950;7:440-3.
3. Nagashima C, Nakashio K, Fujino T. Meningioma and astrocytoma adjacent in the brain. *J Neurosurg* 1963;20:995-9.
4. Strong AJ, Symon L, MacGregor BJ, O'Neill BP. Coincidental meningioma and glioma. Report of two cases. *J Neurosurg* 1976;45:455-8.
5. Marra A, Ramponi G, Grimaldi G. Simultaneous occurrence of right supratentorial meningioma and glioblastoma multiforme. Case report. *Acta Neurochir (Wien)* 1977;36:83-91.
6. Goyal A, Singh AK, Sinha S, Tatke M, Singh D, Gupta V, *et al.* Simultaneous occurrence of meningioma and glioma in brain: Report of two cases. *J Clin Neurosci* 2003;10:252-4.
7. Nestler U, Schmidinger A, Schulz C, Huegens-Penzel M, Gamerding UA, Koehler A, *et al.* Glioblastoma simultaneously present with meningioma – report of three cases. *Zentralbl Neurochir* 2007;68:145-50.
8. Suzuki K, Momota H, Tonooka A, Noguchi H, Yamamoto K, Wanibuchi M, *et al.* Glioblastoma simultaneously present with adjacent meningioma: Case report and review of the literature. *J Neurooncol* 2010;99:147-53.
9. Chen G, Gao X, Liao Y, Xu B. A glioblastoma adjacent to a meningioma. *Br J Neurosurg* 2010;24:718-9.
10. Shahmohammadi MR, Javadian H, Asgari A, Khoshnoud RJ. Glioblastoma simultaneously present with adjacent meningioma. *Int Clin Neurosci J* 2016;3:54-8.