In vitro effect of diode laser against biofilm of *Aggregatibacter actinomycetemcomitans*

Ponpen Tantivitayakul¹, Supanee Rassameemasmaung², Sununtha Thapanabhiboonsuk²

Correspondence: Dr. Supanee Rassameemasmaung Email: supanee.ras@mahidol.ac.th ¹Department of Oral Microbiology, Faculty of Dentistry, Mahidol University, Bangkok, Thailand, ²Department of Oral Medicine and Periodontology, Faculty of Dentistry, Mahidol University, Bangkok, Thailand

ABSTRACT

Objective: The main objective is to investigate the antibacterial effect of diode laser against *Aggregatibacter* actinomycetemcomitans biofilm. **Materials and Methods:** Biofilms of *A. actinomycetemcomitans* plus *Streptococcus* sanguinis grown on bovine root surfaces were treated with an 810-nm diode laser using a noncontact pulsed mode with a pulse interval and pulse length of 20 ms. Four protocols, that is, one episode of 1.5 or 2.5 W for 30 s and three episodes of 1.5 or 2.5 W for 30 s were tested. No treatment and 0.2% chlorhexidine treatment served as negative and positive controls, respectively. Viable bacterial number was determined by colony counting. **Results:** Treatment with chlorhexidine and all laser protocols except that using single episode of 1.5 W reduced the number of *A. actinomycetemcomitans* in either single-species or dual-species biofilm compared to negative control. A higher percentage of *A. actinomycetemcomitans* reduction was demonstrated after increasing the power output or repeating the irradiation. **Conclusions:** The laser protocols used in this study could reduce the number of viable bacteria but not eradicate *A. actinomycetemcomitans* biofilm.

Key words: Aggregatibacter actinomycetemcomitans, biofilm, diode laser, periodontitis treatment

INTRODUCTION

Mechanical periodontal treatment comprising of scaling and root planing aims to remove bacterial plaque, endotoxins, and calculus from the root surface. This measure, together with oral hygiene control, leads to an improved periodontal condition.^[1] However, scaling and root planing are not effective enough in removing such deposits from inaccessible areas^[2] as well as some bacteria including *Aggregatibacter actinomycetemcomitans*^[3] which can invade into the periodontal tissue.^[4]

Diode laser is widely used in the field of dentistry. The ability of diode laser to eliminate pathogens and remove diseased pocket epithelium makes

 Access this article online

 Quick Response Code:

 Website:

 Www.eurjdent.com

it a potential adjunct to conventional mechanical treatment.^[5,6] At present, the antimicrobial property of adjunctive diode laser has been studied with an inconclusive result.^[5,7-10]

In addition to its efficacy, the safety of diode laser is of great importance. To minimize the tissue damage, high laser energy should be avoided. The irradiation should also be split into multiple short episodes instead of using a single episode of long duration.^[11] Thus, this study aimed to test the antibacterial property of an 810-nm diode laser using different irradiation protocols against biofilm of *A. actinomycetemcomitans*.

For reprints contact: reprints@medknow.com

How to cite this article: Tantivitayakul P, Rassameemasmaung S, Thapanabhiboonsuk S. *In vitro* effect of diode laser against biofilm of *Aggregatibacter actinomycetemcomitans*. Eur J Dent 2018;12:485-90. DOI: 10.4103/ejd.ejd_289_17

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

MATERIALS AND METHODS

The biofilm was developed as previously described^[12,13] with slight modifications.

Bacterial strains and culture conditions

To prepare bacterial inoculums, A. actinomycetemcomitans ATCC 29523 or Streptococcus sanguinis ATCC 10556 was grown in Brain Heart Infusion agar (BHI agar; Difco, USA) in a 5% CO₂ incubator at 37°C. Then, 3–5 colonies of each bacterial species were inoculated in 5 mL of BHI broth (Difco) in a 5% CO₂ incubator at 37°C for 18–24 h. Bacterial suspension was adjusted to 1×10^8 colony-forming units/mL (CFU/mL) using spectrophotometer (Biochrom, UK) at an OD₆₀₀. The suspensions were used as inoculums in the biofilm development.

Tooth specimen preparation

Extracted single-rooted bovine teeth were purchased and kept at -80° C until used. The teeth without gross damage or caries lesions were selected. Residual tissue around the root surface was removed. Each tooth was cut and the crown and apical-third portion of the root were discarded. The cervical to middle-third portion of the root was polished with 800-grit sandpaper. The polished surface was left exposed, while the other surfaces of tooth specimen were embedded in silicone putty material (Coltoflax[®], USA) to a size of 10 mm × 10 mm × 5 mm. After that, the specimens were coated with nail varnish, leaving a window of 4 mm × 4 mm in the polished surface for the bacterial biofilm to form. All specimens were sterilized by ethylene oxide.

Biofilm development

Each sterile tooth specimen was placed in an individual well of a 24-well polystyrene plate (Corning, USA). For the formation of single-species biofilm, 1.5 mL of *A. actinomycetemcomitans* suspension was added into each well. For the formation of dual-species biofilm, 750 μ L of *S. sanguinis* and 750 μ L of *A. actinomycetemcomitans* suspension were mixed and transferred into each well. The plate was then incubated at 37°C in a 5% CO₂ incubator shaker for 48 h. Before the treatment, all specimens were washed four times with sterile normal saline solution (NSS) to remove loosely attached bacteria.

Treatment protocols

The experiment comprised four laser irradiation groups, a chlorhexidine (CHX) group and a no

treatment group. Each experimental group was carried out in triplicate and repeated three times on different occasions.

Laser irradiation

Laser treatment was carried out using a diode laser (LaserSmileTM, BIOLASE, Germany) set to a noncontact pulsed mode at a wavelength of 810 nm with a pulse interval of 20 ms and a pulse length of 20 ms. The laser was delivered through an optical fiber with a diameter of 400 μ m. The tip of the fiber was moved, at 0.5–1 mm away from the biofilm surface, with a sweep motion. There were four different laser protocols:

- Protocol A: A single episode of irradiation using a power output of 1.5 W for 30 s (140.625 J/cm²)
- Protocol B: Three episodes of irradiation using a power output of 1.5 W for 30 s (421.875 J/cm²). Each episode was separated by a 30 s pause
- Protocol C: A single episode of irradiation using a power output of 2.5 W for 30 s (234.375 J/cm²)
- Protocol D: Three episodes of irradiation using a power output of 2.5 W for 30 s (703.125 J/cm²). Each episode was separated by a 30 s pause.

After irradiation, each specimen was removed from the silicone block and transferred to a 15 mL microcentrifuge tube containing 2 mL of sterile NSS. The tubes were vigorously vortexed for 3 min. Tenfold serial dilution of bacterial suspension was performed, and it was then plated out on BHI agar. The number of bacterial colonies was counted and calculated as CFU/mL.

Chlorhexidine treatment

Each specimen was removed from the silicone block and transferred to a 1.5 mL microcentrifuge tube containing 1 mL of 0.2% chlorhexidine digluconate (CHX) solution, and then incubated for 30 s. After incubation, the specimens were washed four times with NSS to remove trace amounts of CHX. The specimens were then prepared for colony counting as previously described.

Scanning electron microscopic study

Each specimen was transferred to desiccators until dried. The specimen was coated with a thin layer of gold alloy (100–300 Å) using a sputter coater and viewed under scanning electron microscope (JSM-6610 LV, JEOL, Tokyo) with an accelerating voltage of 20 kV.

Statistical analysis

The CFU/mL data were log-transformed. To identify differences in the log-transformed data between

groups, Kruskal–Wallis nonparametric one-way analysis of variance was performed. A Mann–Whitney U-test was applied for comparison of differences between two groups. Statistical comparisons were performed using SPSS software version 19 (SPSS, IL, USA). For all analyses, P < 0.05 was considered statistically significant.

RESULTS

Bacterial elimination following the treatments

As shown in Table 1, log CFU/mL of *A. actinomycetemcomitans* in the single-species and dual-species biofilms decreased after laser or CHX treatments. All treatment groups except laser Protocol A significantly reduced the number of bacteria compared to the no treatment group [Table 2]. A higher percentage of *A. actinomycetemcomitans* reduction was seen after an increment of the

power output (Protocol C), after repeating the irradiation (Protocol B) or both (Protocol D) in either single-species or dual-species biofilm. The amount of *A. actinomycetemcomitans* after treatment with laser Protocol D was reduced by approximately, 2 log CFU/mL in the single-species biofilm and by 1 log CFU/mL in the dual-species biofilm. However, CHX treatment exhibited >3 log CFU/mL reduction.

Biofilm morphology from scanning electron microscopy

In a 48 h-old, single-species biofilm, numerous coccobacilli were found on the bovine root surface [Figure 1a]. In a 48 h-old, dual-species biofilm, the aggregation of cocci and coccobacilli were found. Some formed a cluster of bacteria and some formed layers on top of each other [Figure 1b].

After laser treatment, the mass of extracellular matrix covering bacteria was found on the root

Table 1: Mean±standard deviation of log colony-forming units/mL and percentage of bacterial reduction following treatments

Group	Mean±SD of log CFU/mL (percentage reduction [#])							
	A. actinomycetemcomitans in SSB (%)	A. actinomycetemcomitans in DSB (%)	Total bacteria (%)					
No treatment	7.66±0.51	7.16±0.19	7.49±0.33					
A	7.57±0.40 (41.739)	6.89±0.46 (23.387)	7.37±0.42 (9.820)					
В	6.12±0.26 (98.212)	6.08±0.27 (91.210)	6.50±0.34 (89.524)					
С	6.53±0.41 (94.411)	6.40±0.32 (80.645)	6.73±0.26 (84.288)					
D	5.80±0.36 (99.000)	5.94±0.51 (90.645)	6.22±0.50 (93.425)					
CHX	4.40±0.82 (99.902)	1.94±0.48 (99.999)	2.92±0.71 (99.994)					

*Percentage reduction between no treatment and treatment groups. Groups - A: 1.5 W; 30 s × one episode, B: 1.5 W; 30 s × three episodes, C: 2.5 W; 30 s × one episode, D: 2.5 W; 30 s × three episodes. CHX: 0.2% chlorhexidine digluconate 30 s, *A. actinomycetemcomitans*: *Aggregatibacter actinomycetemcomitans*, DSB: Dual-species biofilm, SSB: Single-species biofilm, CFU: Colony-forming units, SD: Standard deviation

Table 2: P values for the differences in the log colony-forming units/mL of Aggregatibacter actinomycetemcomitans and total bacteria following treatments

	-					
	Group	A	В	С	D	СНХ
A. actinomycetemcomitans in SSB	No treatment	0.82	<0.001*	<0.001*	<0.001*	<0.001*
	A		<0.001*	<0.001*	<0.001*	<0.001*
	В			0.031*	0.063	<0.001*
	С				0.002*	<0.001*
	D					<0.001*
A. actinomycetemcomitans in DSB	No treatment	0.486	0.029*	0.029*	0.029*	0.029*
	А		0.029*	0.114	0.057	0.029*
	В			0.2	0.686	0.029*
	С				0.2	0.029*
	D					0.029*
Total bacteria (DSB)	No treatment	0.433	<0.001*	<0.001*	<0.001*	<0.001*
	А		<0.001*	0.001*	<0.001*	<0.001*
	В			0.136	0.297	<0.001*
	С				0.024*	<0.001*
	D					<0.001*

*Statistically significant difference using Mann-Whitney U-test. Groups - A: 1.5 W; 30 s × one episode, B: 1.5 W; 30 s × three episodes, C: 2.5 W; 30 s × one episode, D: 2.5 W; 30 s × three episodes. CHX: 0.2% chlorhexidine digluconate 30 s, *A. actinomycetemcomitans: Aggregatibacter actinomycetemcomitans*, DSB: Dual-species biofilm, SSB: Single-species biofilm



Figure 1: Scanning electron micrographs of (a) single-species biofilm (original magnification × 7000) and (b) dual-species biofilm (×5000) on bovine root surface

surface, while the morphology of a few bacteria with irregular shape was observed in the dual-species biofilm. Few microbes remained on the root surface [Figure 2a and b].

DISCUSSION

Mechanical debridement is a standard method for periodontal therapy because it is a cause-related approach. This mode of treatment results in a significant reduction of several periodontal pathogens including Porphyromonas gingivalis and Tannerella forsythia. However, the level of A. actinomycetemcomitans does not significantly decrease after therapy.^[3] A. actinomycetemcomitans is known to involve in the inflammation-driven alveolar bone loss through regulation of chemokine signaling in several cell types.^[14] Its presence is considered a risk marker for the progression of attachment loss.^[15] Since the elimination of bacteria is important to delay subgingival recolonization, many approaches including the use of antibiotics^[16] or antimicrobial photodynamic therapy (PDT)^[17,18] have been introduced. However, these bacteria can resist such therapeutic procedures as shown by the reduced antibiotic susceptibility.^[16] Regarding PDT, not all studies could decrease bacteria by at least 3 log CFU/mL (99.9%).[17]

When reviewing the antimicrobial effect of diode laser, conflicting results were obtained, partly because of differences in the irradiation parameters used.^[7-10] For example, Moritz *et al.*^[8] found that three episodes of irradiation (at 1 week, 2 months, and 4 months after scaling and root planing) with an 805-nm diode laser, set at 2.5 W with a pulse duration of 10 ms and a pulse rate of 50 Hz, led to a significant bacterial reduction at 6 months over H_2O_2 rinsing of the periodontal pockets. In contrast, Song *et al.*^[10] reported that two episodes of irradiation with an 810-nm diode laser using a power of 0.8 W in a continuous-wave mode for 30 s could kill only 17%–54% of the multispecies biofilm. Kreisler *et al.*^[7] investigated the antimicrobial effect of



Figure 2: Scanning electron micrographs of (a) single-species biofilm and (b) dual-species biofilm (×5000) after laser treatment using Protocol D

an 809-nm diode laser against *S. sanguinis* on dental implant surfaces using a power output between 0.5 W and 2.5 W and found that bacteria could be reduced by approximately 50% at 0.5 W to 99% at 2.5 W.

In terms of thermal damage, the critical temperature that can induce heat injury varies between tissues. Kreisler et al.^[19] evaluated the morphologic alterations of the root surface after a noncontact 809-nm GaAlAs laser irradiation. They found no alterations in lasing dry or saline-moistened root specimens. However, blood-coated specimens showed partial or total carbonization after irradiation with ≥1.5 W in a continuous mode for 10-30 s. Kreisler et al.[11] also investigated intrapulpal temperature change during root surface irradiation with an 809-nm GaAlAs laser. With a remaining dentin thickness of 1 mm, only irradiation at 0.5 W for a maximum of 10 s did not increase an intrapulpal temperature beyond a critical threshold. When irradiation was stopped for 30-40 s, the temperature returned to baseline. These authors suggested that irradiation should be interrupted to allow the hard tissue and pulp to cool down.

According to the manufacturer's recommendation, the diode laser parameters for treating periodontal pockets are 1.5 W in a pulsed mode with a pulse interval of 20 ms and a pulse length of 20 ms. The recommended duration is 30–240 s depending on the severity of inflammation. Thus, these parameters were chosen in this study. In addition, we questioned whether the sterilization effect would be increased if the irradiation was delivered three times with a 30 s pause between each episode. The reason for this was to increase the laser energy while allowing the elevated temperature to return to baseline.^[11]

The bactericidal mechanism of a diode laser is primarily based on the photothermal effect. Pirnat*etal*.^[18] indicated that bacterial cell death is induced by short-term localized heating of bacterial microenvironment to a lethal temperature. In this study, the reduction of *A. actinomycetemcomitans* after laser treatment of 1.5 W

for 30 s was not significantly different compared to negative control. However, laser irradiation of 1.5 W for 30 s, when performed three times, significantly decreased *A. actinomycetemcomitans* compared to the groups receiving a single episode of irradiation. The killing effect of diode laser could result from heat denaturation of macromolecules. Another possible killing mechanism might result from the endogenous photosensitizers of *A. actinomycetemcomitans* which can be light-activated and generate singlet oxygen.^[20] In this study, no attempt was made to find the mechanism of bacterial killing.

When the power output was increased to 2.5 W, a lower number of bacteria was found. Furthermore, the log CFU/mL of A. actinomycetemcomitans in the single-species biofilm was significantly different after irradiation at 2.5 W for three times, compared to that irradiated with a single episode of 2.5 W (P = 0.002). However, a borderline nonsignificant difference (P = 0.057) in the log CFU/mL of A. actinomycetemcomitans in the dual-species biofilm was found between the group receiving 1.5 W for 30 s and that receiving three episodes of 2.5 W for 30 s, which might be explained by a small number of samples. Thus, diode laser could decrease the number of A. actinomycetemcomitans in the biofilm form. However, if a minimum of 99.9% bacterial reduction must be achieved to claim an antibacterial effect, no laser protocols used in this study showed antibacterial activity against biofilm of *A. actinomycetemcomitans*.

A dual-species biofilm of *A. actinomycetemcomitans* and *S. sanguinis* was used in this study to observe the effect of the diode laser against mixed-species bacterial biofilm. *S. sanguinis* is the most prominent bacterial species among the primary colonizers of dental plaque.^[21] Therefore, *S. sanguinis* acts as a bridge between the tooth surface and *A. actinomycetemcomitans*. In this study, the percentage of bacterial reduction after laser irradiation was less in the dual-species biofilm than in the single-species biofilm, suggesting more resistance of dual-species biofilm to laser therapy. This can be explained by a more complex set of polysaccharides or a thicker biofilm produced by more species of bacteria.

PDT has been used for an antimicrobial purpose in dentistry.^[22] However, the results vary depending on the type and concentration of photosensitizer as well as the incubation time for the photosensitizer to reach its target.^[17] Mattiello *et al.*^[23] studied an *in vitro* photodynamic effect using 0.01% toluidine blue-O

for 5 min and an AlGaInP diode laser for 3 min on *A. actinomycetemcomitans* and *S. sanguinis* inoculums and found a 61.53% reduction of the CFU counts for *A. actinomycetemcomitans*. Compared to that study, our results showed a greater reduction of CFU count for *A. actinomycetemcomitans* after irradiation using one episode of 2.5 W for 30 s or three episodes of 1.5 W for 30 s.

In this study, the *in vitro* laser application could reduce clinical variations such as pocket depth or presence of bleeding. However, the use of a single-species or dual-species biofilm was considered a limitation because it could not mimic the polymicrobial biofilm found in the oral cavity.

CONCLUSIONS

The laser protocols used in this study could reduce the number of viable bacteria but could not eradicate the biofilm of *A. actinomycetemcomitans*.

Acknowledgment

We would like to thank Assistant Professor Chulaluk Komoltri for statistical analysis, Miss Chayada Teanchai for preparing SEM specimens, and Miss Thaniya Muadchiangka for microbiological laboratory.

Financial support and sponsorship

This study was financially supported by Mahidol University Faculty of Dentistry Grant (04/2557).

Conflicts of interest

There are no conflicts of interest.

REFERENCES

- Badersten A, Nilveus R, Egelberg J. Effect of nonsurgical periodontal therapy. II. Severely advanced periodontitis. J Clin Periodontol 1984;11:63-76.
- Parashis AO, Anagnou-Vareltzides A, Demetriou N. Calculus removal from multirooted teeth with and without surgical access. (I). Efficacy on external and furcation surfaces in relation to probing depth. J Clin Periodontol 1993;20:63-8.
- Takamatsu N, Yano K, He T, Umeda M, Ishikawa I. Effect of initial periodontal therapy on the frequency of detecting *Bacteroides forsythus*, *Porphyromonas gingivalis*, and *Actinobacillus actinomycetemcomitans*. J Periodontol 1999;70:574-80.
- Saglie FR, Smith CT, Newman MG, Carranza FA Jr., Pertuiset JH, Cheng L, *et al.* The presence of bacteria in the oral epithelium in periodontal disease. II. Immunohistochemical identification of bacteria. J Periodontol 1986;57:492-500.
- 5. Mizutani K, Aoki A, Coluzzi D, Yukna R, Wang CY, Pavlic V, *et al.* Lasers in minimally invasive periodontal and peri-implant therapy. Periodontol 2000 2016;71:185-212.
- 6. Pirnat S. Versatility of an 810 nm diode laser in dentistry: An overview. J Laser Health Acad 2007;4:1-9.
- Kreisler M, Kohnen W, Marinello C, Schoof J, Langnau E, Jansen B, et al. Antimicrobial efficacy of semiconductor laser irradiation on implant surfaces. Int J Oral Maxillofac Implants 2003;18:706-11.

- Moritz A, Schoop U, Goharkhay K, Schauer P, Doertbudak O, Wernisch J, *et al.* Treatment of periodontal pockets with a diode laser. Lasers Surg Med 1998;22:302-11.
- Sennhenn-Kirchner S, Klaue S, Wolff N, Mergeryan H, Borg von Zepelin M, Jacobs HG, *et al.* Decontamination of rough titanium surfaces with diode lasers: Microbiological findings on *in vivo* grown biofilms. Clin Oral Implants Res 2007;18:126-32.
 Song X, Yaskell T, Klepac-Ceraj V, Lynch MC, Soukos NS.
- Song X, Yaskell T, Klepac-Ceraj V, Lynch MC, Soukos NS. Antimicrobial action of minocycline microspheres versus 810-nm diode laser on human dental plaque microcosm biofilms. J Periodontol 2014;85:335-42.
- Kreisler M, Al-Haj H, D'Hoedt B. Intrapulpal temperature changes during root surface irradiation with an 809-nm gaAlAs laser. Oral Surg Oral Med Oral Pathol Oral Radiol Endod 2002;93:730-5.
- 12. Folwaczny M, Mehl A, Aggstaller H, Hickel R. Antimicrobial effects of 2.94 microm er: YAG laser radiation on root surfaces: An *in vitro* study. J Clin Periodontol 2002;29:73-8.
- 13. Walker C, Sedlacek MJ. An *in vitro* biofilm model of subgingival plaque. Oral Microbiol Immunol 2007;22:152-61.
- Herbert BA, Novince CM, Kirkwood KL. Aggregatibacter actinomycetemcomitans, a potent immunoregulator of the periodontal host defense system and alveolar bone homeostasis. Mol Oral Microbiol 2016;31:207-27.
- 15. Van der Velden U, Abbas F, Armand S, Loos BG, Timmerman MF, Van der Weijden GA, *et al.* Java project on periodontal diseases. The natural development of periodontitis: Risk factors, risk predictors and

risk determinants. J Clin Periodontol 2006;33:540-8.

- Lewis K. Riddle of biofilm resistance. Antimicrob Agents Chemother 2001;45:999-1007.
- Haag PA, Steiger-Ronay V, Schmidlin PR. The *in vitro* antimicrobial efficacy of PDT against periodontopathogenic bacteria. Int J Mol Sci 2015;16:27327-38.
- Pirnat S, Lukac M, Ihan A. Study of the direct bactericidal effect of nd: YAG and diode laser parameters used in endodontics on pigmented and nonpigmented bacteria. Lasers Med Sci 2011;26:755-61.
- Kreisler M, Al Haj H, Daubländer M, Götz H, Duschner H, Willershausen B, *et al*. Effect of diode laser irradiation on root surfaces *in vitro*. J Clin Laser Med Surg 2002;20:63-9.
- Cieplik F, Späth A, Leibl C, Gollmer A, Regensburger J, Tabenski L, et al. Blue light kills Aggregatibacter actinomycetemcomitans due to its endogenous photosensitizers. Clin Oral Investig 2014;18:1763-9.
- Kolenbrander PE, Ganeshkumar N, Cassels FJ, Hughes CV. Coaggregation: Specific adherence among human oral plaque bacteria. FASEB J 1993;7:406-13.
- Ricatto LG, Conrado LA, Turssi CP, França FM, Basting RT, Amaral FL, et al. Comparative evaluation of photodynamic therapy using LASER or light emitting diode on cariogenic bacteria: An *in vitro* study. Eur J Dent 2014;8:509-14.
- Mattiello FD, Coelho AA, Martins OP, Mattiello RD, Ferrão Júnior JP. In vitro effect of photodynamic therapy on Aggregatibacter actinomycetemcomitans and Streptococcus sanguinis. Braz Dent J 2011;22:398-403.