

Author's Reply

Sir,

We agree with the pertinent remarks of the authors and appreciate their interest in our article. Differentiating tuberculosis from sarcoidosis with absolute certainty is difficult because of overlapping clinical and radiological features.^[1] In such cases, tissue sampling is usually done; nevertheless, this distinction may not be achieved in all cases despite sampling. Even with the addition of microbiological investigations, the diagnostic yield is at best around 60%.^[2-5]

The purpose of this review article was to highlight the imaging similarities and differentiating features between the two entities with only a brief description of clinical and laboratory correlates. We did mention in the imaging-based algorithm proposed in our article about the use of microbiological test—CBNAAT (Gene Xpert) in initial sputum evaluation which is a part of RNTCP guidelines for assessment of tuberculosis.^[6] In general, the investigation is routinely performed on any specimen obtained in a suspected case of tuberculosis, for example, bronchoalveolar lavage/fine needle aspiration/biopsy specimens. However,

discussion of the implications of it being positive or negative in correlation with the pathological presence or absence of non-caseating granulomas is beyond the range of an imaging-based discussion on sarcoidosis/tuberculosis. Microbiological and pathological evaluation of this spectrum of granulomatous diseases is a complex topic in itself and requires a thorough search and analysis of the literature. Hence, inclusion of the same is beyond the purview and intent of the current article.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

**Ashu S Bhalla, A Das, P Naranje, A Goyal,
R Guleria¹, Gopi C Khilnani¹**

Departments of Radiodiagnosis, ¹Pulmonary Medicine and
Sleep Disorders, All India Institute of Medical Sciences,
New Delhi, India
E-mail: ashubhalla1@yahoo.com

References

1. Bhalla AS, Das A, Naranje P, Goyal A, Guleria R, Khilnani GC. Dilemma of diagnosing thoracic sarcoidosis in tuberculosis endemic regions: An imaging-based approach. Part 1. *Indian J Radiol Imaging* 2017;27:369-79.
2. Thangakunam B, Isaac BTJ, Christopher DJ. Endobronchial ultrasound experience in a high tuberculosis prevalence setting. *Indian J Tuberc* 2017;64:196-200.
3. Eom JS, Mok JH, Lee MK, Lee K, Kim MJ, Jang SM, *et al.* Efficacy of TB-PCR using EBUS-TBNA samples in patients with intrathoracic granulomatous lymphadenopathy. *BMC Pulm Med* 2015;15:166.
4. Prasoon D. Acid-fast bacilli in fine needle aspiration smears from tuberculous lymph nodes. Where to look for them. *ActaCytol* 2000;44:297-300.
5. Bezabih M, Mariam DW, Selassie SG. Fine needle aspiration cytology of suspected tuberculous lymphadenitis. *Cytopathol* 2002;13:284-90.
6. Bhalla AS, Das A, Naranje P, Goyal A, Guleria R, Khilnani GC. Dilemma of diagnosing thoracic sarcoidosis in tuberculosis endemic regions: An imaging-based approach. Part 2. *Indian J Radiol Imaging* 2017;27:380-8.

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

Access this article online	
Quick Response Code:	Website: www.ijri.org
	DOI: 10.4103/ijri.IJRI_85_18

Cite this article as: Bhalla AS, Das A, Naranje P, Goyal A, Guleria R, Khilnani GC. Author's Reply. *Indian J Radiol Imaging* 2018;28:268-9.

© 2018 Indian Journal of Radiology and Imaging | Published by Wolters Kluwer - Medknow