

## Letter to the Editor

### Saree cancer: A case report

DOI: 10.4103/sajc.sajc\_16\_16

Dermatoses are more commonly seen in body folds and in waist areas where sari is tightened. We are presenting a rare case of saree cancer in the waistline in an 80-year-old elderly female.

An 80-year-old female presented with a long-standing swelling with oozing in the right waist. Waistline showed an ulceroproliferative cauliflower like growth measuring 8 cm × 7 cm. [Figure 1] was seen with hyper- and hypo-pigmented patches surrounding the lesion. Bilateral inguinal lymphadenopathy was present. Biopsy of growth showed well-differentiated squamous cell carcinoma (SCC) [Figure 2]. The patient gave a history of wearing saree for 70 years.

Khanolkar and Suryabai<sup>[1]</sup> described “dhoti cancer” in Indian males. A similar type of “saree cancer” in females was described by Patil *et al.*<sup>[2]</sup>

Pigmentation and mild scaling over the waist in Indian females who wear saree are so common that they consider it normal and ignore it.

The incidence of malignancy in scar tissues is 0.1%–2.5%. The malignant change in the form of SCC is seen in Marjolin’s ulcer.

Wide local excision of tumor with inguinal block dissection is the treatment of choice while the combination of surgery, radiotherapy, and chemotherapy may be of extra value compared to surgery alone.<sup>[3]</sup>



Figure 1: Growth along right side of waist and hypopigmentation on the left side




Figure 2: Well-differentiated squamous cell carcinoma (H and E, 100)

(Continue on page 107...)

(Letter to the editor continue from page 101...)

SCCs which develop on chronic skin lesions have a higher incidence of metastasis (9%–36%) as compared to those arise in the previously normal skin (1%–10%). Bilateral inguinal lymph nodes were enlarged in our case. Fine-needle aspiration or biopsy of inguinal lymph nodes was not done in our case; hence, we cannot comment about metastasis.

Pigmentation and mild scaling over the waist are so common in Indian females that they consider it normal and ignore it. This case is presented for its rarity and to bring awareness about saree cancer among Indian women to detect it earlier.

#### Declaration of patient consent

The authors certify that they have obtained all appropriate patient consent forms. In the form the patient(s) has/have given his/her/their consent for his/her/their images and other clinical information to be reported in the journal. The patients understand that their names and initials will not be published and due efforts will be made to conceal their identity, but anonymity cannot be guaranteed.

#### Financial support and sponsorship

Nil.

#### Conflicts of interest

There are no conflicts of interest.

**Chitrawati Bal Gargade, Abhay Yashwant Desai**

Department of Pathology, BKL Walawalkar Rural Medical College, Chiplun, Maharashtra, India

Correspondence to: Dr. Chitrawati Bal Gargade,  
E-mail: chitragargade@yahoo.com

#### References

1. Khanolkar VR, Suryabai B. Cancer in relation to usages; three new types in India. Arch Pathol (Chic) 1945;40:351-61.
2. Patil AS, Bakhshi GD, Puri YS, Gedham MC, Naik AV, Joshi RK. Saree cancer. Bombay Hosp J 2005;47:302-3.
3. Takalkar UV, Asegaonkar SB, Kodlikeri P, Kulkarni U, Borundiya V, Advani SH. Saree cancer in Indian woman treated successfully with multimodality management. Dermatol Reports 2014;6:5128.

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

**How to cite this article:** Gargade CB, Desai AY. Saree cancer: A case report. South Asian J Cancer 2019;8:101-7.  
© 2019 The South Asian Journal of Cancer | Published by Wolters Kluwer - Medknow