Accelerating the Benefits of the Problem Oriented Medical Record

Joel Buchanan¹

¹Department of Medicine, University of Wisconsin Madison School of Medicine and Public Health, Madison, Wisconsin, United States

Correspondence to:

Joel Buchanan, MD UW Health Information Services 8007 Excelsior Drive Madison, WI 53717 Email:jbuchanan@uwhealth.org

Appl Clin Inform 2017; 8: 180–190

https://doi.org/10.4338/ACI-2016-04-IE-0054 received: April 10, 2016 accepted: December 11, 2016 published: February 15, 2017 **Citation:** Buchanan J. Accelerating the benefits of the problem oriented medical record. Appl Clin Inform 2017; 8: 180–190 https://doi.org/10.4338/ACI-2016-04-IE-0054

Background

In 1968, Weed wrote the seminal article on the Problem Oriented Medical Record (POMR), *Medical Records that Guide and Teach* [1]. He promoted two major concepts: the Problem Oriented Medical Record framework and a particular *note* structure. Weed wrote that a system with records organized by problems should replace systems organized by source and systems which lacked an organizing framework (both common practices in 1968). He suggested that progress notes should adhere to a structure with five elements: Subjective, Objective, Rx, Interpretation and Plan [1, 2]. This five-component structure evolved to the four-component Subjective, Objective, Assessment and Plan structure, now commonly known as SOAP. Although the SOAP note structure has been widely embraced, adoption of the POMR has been more limited.

Given the numerous potential benefits, Simons et al. recently surveyed the literature and published the attributes (called *determinants*) necessary for success of the POMR [3]. The authors list twelve determinants, including functionality, completeness, and efficiency. One aspect of functionality is specified as "context dependent views of all data". A framework using these context (problem) dependent views is the *Problem Oriented View* (POV). The use of POV to accelerate POMR adoption is the focus of this editorial.

This editorial will highlight some crucial studies and commentary on the POMR, outline a POV model and discuss the use of problem concept maps (PCMs) as an enabling methodology to generate the POV within an EHR. The POV will save busy clinicians time and work, as the computer will be able to organize and display data by problem, while continuing to offer current display frameworks.

The Meaningful Use Stage 1 Core Measure specification defines the problem list as *a list of current and active diagnoses as well as past diagnoses relevant to the current care of the patient* [4]. Clinician attitudes about using the problem list vary widely [5]. Problem lists are subject to the network effect, in which goods or services becomes more valuable when more people use them [6]. Because problem lists have not been fully adopted, those of us who do use the framework do not experience the maximum potential benefit. Investigators have addressed the challenge of incomplete adoption by working on tools to improve completeness [7, 8, 9] and by suggesting approaches for lowering the complexity of problem list maintenance [10, 11].

The advent of the EHR seemed to offer a path toward implementation of the POMR. Even in 1968, one of the many striking aspects of Weed's article was his vision for using the computer to leverage his concepts. However, the progress on this path has been laborious and slow. Strict adherence to Weed's recommendations involves creating formatted problem-oriented progress notes. The use of such notes likely helps to improve clinical thinking, especially among trainees [12, 13]. In theory, charting by creating problem-oriented progress notes on a computer allows others to view an organized document sorted by problem. However, many EHR users do not generate problem-oriented notes because the creation tools are so inefficient. The POV model will allow EHR users to see data and documentation organized into a problem-oriented view, *even if prior notes were not created using computerized problem-oriented charting*.

For many years, EHRs have been organized primarily by data source (clinic notes, hospitalizations, lab results, imaging reports, diagnostic procedure, etc.) and secondarily by chronologic or reverse chronologic order within that data source. Separate storage and indexing of inpatient and outpatient information tend to fragment individual patient records. The EHR offers the tantalizing possibility of adding a POMR view to the established source/chronologic-oriented medical record. However, the realization of a computerized problem-oriented view has been quite challenging for vendors and users.

Paper records lent themselves to a chronologic or reverse chronologic structure. Hospitalizations were marked with tabs that showed the admission and discharge dates (> Figure 1). Lab and x-ray reports were placed in the chart using the *shingle method* (> Figure 2).

As they created computer interfaces for data retrieval, many EHR vendors "paved the cowpath" [14] when they computerized the chronologically oriented scheme of the paper record. Within each domain (notes, lab data, medications, imaging, and diagnostic test results), vendors created an electronic chronologic index. This type of orientation can be useful, especially when the user needs to focus on the most recent data in a domain. However, a display with aggregated data [15] is essential

J Buchanan. Accelerating the benefits of the problem oriented medical record

to realize Dr. Weed's vision of the problem-oriented record. Vendors should give users both options, providing an EHR that allows a viewer to toggle back and forth easily between a chronologic and a problem-based display.

Without computerized data aggregation, EHR users who wish to view data by problem must organize the data themselves. They must establish links from the problems to relevant data and notes, and/or copy and paste lab, imaging, and procedure data and consultant notes into the annotation section of the problem list. This is a time-consuming task for users, and one that is not practiced consistently.

Model Approach

A user-friendly problem based record should offer on-demand display of aggregated data and notes relevant to a particular problem. These data or links to these data will only display when the user selects the problem. When that problem is not being reviewed, the data display will collapse and take up no screen space.

By combining the problem list with problem concept maps (PCMs), our model generates a dynamic display of the POMR. PCMs are similar to problem concept-oriented views, which are well described by other investigators [16,17]. We propose using the problem list and PCMs to automatically create a POV. The user initially sees the problem list (> Figure 3). When the user clicks on the problem of interest, the system creates the POV for that problem (> Figure 4). This view is dynamically created *at the time of display, rather than trying to store the data in a problem-based structure.*

Model Implementation Methodology

For common problems, hard coded PCMs will be stored at the system level. The concept maps will be built with standard public domain vocabularies such as SNOMED for problems, RxNorm for medications, LOINC for laboratory results and radiology results, and ICD-10 codes for encounters/ hospitalizations. The vocabulary for diagnostic procedures remains to be determined. The EHR will use these PCMs to display medications, results (lab, imaging, and diagnostic procedures), and relevant documentation in a problem-oriented manner. The first component of a map is a cluster of SNOMED codes, which represent the problems that will trigger the system to use the map. Then relevant medications and lab tests will be added to the map, using RxNorm codes and LOINC codes, respectively. Additional codes for imaging and diagnostic procedures will be added. Finally visit diagnosis codes are added. The medication and lab section of a diabetes problem concept map is shown in ▶ Figure 5.

The individual *map component links* that constitute Problem Concept Maps will be used, along with *indication links* and *manual links*, to create the dynamic displays needed for the problemoriented view. These three link types are defined in ▶ Table 1. Although providers often do not currently document the indication for a prescribed medication, this should change as providers see the numerous advantages of recording the indication [18].

► Table 2 shows the functionality, maps and links needed for each data type to generate a robust dynamic POV. PCMs containing component links are the building blocks of our model, enabling the computer to create a dynamic display of aggregate data.

In this model, the system can use concept maps to create a dynamic display based on *relationships drawn from the maps at the time the display is created*. If no PCM is available, the system can use any existing manual links or indication links to create a dynamic display based on *relationships previously created* for that patient's record. Though the use of indication links is likely to grow (see above), the burden of creating manual links will be relieved as PCMs and automatic dynamic displays become more widespread.

Various vendors provide some of the items listed in \triangleright Table 2. At the University of Wisconsin, Epic provides the functionality for items 1, 2, 3, 4 and 5, and UW has implemented concept maps for medications and labs, for seven problems.

This document was downloaded for personal use only. Unauthorized distribution is strictly prohibited.

Model POV Example

Suppose that the provider views the problem list (Figure 3), focuses on a particular problem and selects it. Using the SNOMED code for that problem, the system retrieves the associated PCM. The system then uses the RxNorm, LOINC, and other codes in that PCM to find relevant data in the individual patient record containing these codes. The data retrieved are shown on the screen, and aggregated under that particular problem (Figure 4). A problem concept map will also contain the provider specialties, clinic specialties and billing diagnoses related to that particular problem. The system can then use the map with scheduling and billing codes to identify relevant encounters and display links to the documentation from those encounters.

For a given problem, POV software will display links to entire notes related to the associated billing diagnosis or specialty (\triangleright Figure 6). Additional specificity will be gained if the note author has documented the name of the problem just before the assessment and plan paragraphs for that problem. Then the system can use simple natural language processing to link directly to that section of the assessment and plan (\triangleright Figure 7). The use of such a direct link will save time for users.

Additional Considerations

The Problem Oriented View is intended for viewing and to assist with clinical decision making. POV is not intended for note generation. Users may be tempted to use POV to create progress notes that serve as all inclusive summaries. With a few exceptions such as consult notes, this temptation should be avoided, because it will accelerate note bloat. Instead, users should develop confidence that they can use POV to generate a problem-based aggregate display at any time, rather than searching for prior summary notes that are created repeatedly at each encounter.

The title of a given problem can change over time, as when an *initial disease* evolves into a different problem (e.g., *impaired fasting glucose* evolves to *diabetes*), or a *symptom problem* is identified to be a *disease process* (e.g., *shortness of breath* is found to be caused by *chronic pulmonary emboli*). For problem-data relationships based on problem concept maps, these title changes will cause no difficulty, because the system will dynamically generate the data display when the user clicks on the linked new problem name. For problem-data relationships based on an indication links or other type of links, the physician may need to generate links to the new problem title. Then an accurate problem-oriented view will be accessible.

We have created five problem concept maps using expert consensus across six academic medical centers. This process combines a modified Delphi technique with consensus decision making. The experts first vote and make suggestions anonymously. Then they interact asynchronously using Google Drive to advance toward consensus. In the last step, the consensus is finalized in a one hour conference call. The time needed from each volunteer subject matter expert is 2 to 4 hours. Paid staff perform central coordination of the process. We will be posting maps at problemlist.org.

Success of the Problem Oriented View model will require significant curation and knowledge management resources for construction and maintenance of the Problem Concept Maps. Just as LOINC and SNOMED undergo periodic review [19,20], each of the problem concept maps will need yearly review. As the RxNorm, LOINC, and SNOMED vocabularies require paid professional staff to provide long term support, the problem concept maps will require paid staff to facilitate the work of the volunteer subject matter experts.

Summary

The Problem Oriented View model we have described will generate an on-demand dynamic display of aggregated data and documentation (or index hyperlinks to those data and documentation) in the problem list. The existence of a Problem Concept Map along with a problem list entry for the problem of interest will allow the user to easily view aggregate data and documentation for that problem. The original data and documentation will continue to be stored in the usual manner. Existing reverse chronologic views of data will still be available. Implementation of this model will bring us significantly closer to Weed's vision of the Problem Oriented Medical Record.

Multiple Choice Questions

1. What is the network effect?

- A The phenomenon by which the availability of multiple wireless access points in a building allows better access to the Internet
- B The problem in which too many people using the same wireless access point causes overcrowding and decreased Internet speed
- C The phenomenon by which goods or services become more valuable when more people use them
- D The problem in which too many wireless access points in the same building cause interference and corruption of Internet traffic

The correct answer is (C). The *network effect* is the effect that one user of goods or services has on the value of those products to other people. When a network effect is present, the value of a product or service is dependent on the number of others using it. [6,21] Classic examples of the network effect include the telephone and the Internet.

If too many people are using the same wireless access point or too many wireless access points in a building cause interference, then there is said to be *network congestion*. The availability of multiple wireless access points may indeed allow better access to the Internet, but this occurs via relief of network congestion and is *not* called the network effect.

2. The Meaningful Use Stage 1 Core Measure specification defines the problem list as:

- A A list of all past illnesses, all past surgeries, and ongoing medical conditions
- B A list of current and active diagnoses as well as past diagnoses relevant to the current care of the patient
- C A list first described by Dr. William Osler in 1868, and which facilitates the creation of a Problem Oriented Medical Record
- D A list of medical conditions that require ongoing treatment and monitoring, for example diabetes

The correct answer is (B). The Meaningful Use Stage 1 Core Measure specification defines the problem list as a list of current and active diagnoses as well as past diagnoses relevant to the current care of the patient. [4]

The specification makes no explicit mention of including past surgeries, although providers may choose to include previous surgeries relevant to current care. There is no requirement to include all past illnesses. The Problem Oriented Record and problem list were described in 1968 by Dr. Lawrence Weed.

Conflicts of Interest

The author has no conflicts of interest to report.

Human Subjects Research Statement

This manuscript did not involve research on human subjects

Acknowledgements

Thanks to DuWayne Willett, Richard Bruce, Eneida Mendonça, Scott Springman, Deborah Dunham, Julie Fagan and Jeff Alexander for review and comments.




Fig. 1 Record of Hospital Admissions in Chronological Order

	MUPORT AN IN		
	REPORT IN ME		
	() INPATIENT		
D.	COPD GROUP	DIAGNOST ONE REQUEST PER CONSU EXAM REQUESTED:	IC RADIOLOGY
Ø.		ift Ler 1-51	here
10.5		SALIENT CLINICAL DATA - PP	OVISION DIAGNOSIS
*Xes		Fall last na	STAT
18-000 77			
POSSIBLY PREGNANT YES () NO ()	BEQUESTED BY	DATE REQUEST	DATE EXAM
ARRIVAL: AMB () WHEEL CH () CART ()		18 EMERGENCY RC nh Dec 77	18 Dec. 77 DOM DAY NO.
AP AND LATERAL SPINE: There is a contrast material present.	narrowing of L5 on S	1. No evidence of fractu	re. Previous
ER 152406			
		trans (R)	22
UNRI 630 (1078)	UNIVERSITY OF WISCONSIN	12.	MEDICAL CHART
JWH 610 (1076)			
	UNIVERSITY OF WISCONSIN	HOSPITALS	MEDICAL CHART
AN 870 (10776)	INIVERSITY OF WISCONSIN	HOSPITALS	MEDICAL CHART
(WH \$10 (10/76)	UNIVERSITY OF WISCONSIN	HOSPITALS	MEDICAL CHART
	X-RAY REPO	RTS	
		annen an	

Fig. 2 X-Ray Reports Stored with the Shingle Method

Applied Clinical Informatics 186

Problem List	
Endocrine	
Diabetes	
Musculoskeletal	
Degenerative Joint Disease	
Neurologic	
Epilepsy	

Fig. 3 Problem List without Showing any Aggregated Data

Problem List							
Endocrine							
Diabetes							
	Musculoskeletal						
D	Degenerative Joi	nt Disease					
Neu	rologic						
\bigtriangledown	Epilepsy						
	Meds						
	lamOTRIGINE (LAMICTAL) 100MG tab	Take 2 tabs (200m) 1.5 tabs (150mg) ir per G-Tube)	g) in the AM & 1 the PM. (Take crushed	105 tab	6 ordered	6/10/2015	
	midazolam (VERSED) 2MG/ML syrup	· · · · · · · · · · · · · · · · · · ·				7/2/2015	
	Labs						
	LAMOTRIGINE	4.8 1/11/2014					
	Imaging						
	9/12/2012	MRI HEAD W & W/O CONTRAST					
	Impression: 1. Find	dings compatible with bilateral Sturge-Weber syndrome with associated					
	Procedures						
	12/14/2012	Routine EEG					
	CLINICAL INTERPRETATION: This EEG recording is abnormal due to generalized slowing and disorganization.						
	Clinic Notes						
	3/11/2015	Epilepsy Dr. Stanley					
	9/23/2014 Neurosurgery Dr. Livingstone						
Hospitalizations							
	7/31/2014	Neurosurgery	Dr. Livingstone				
	8/2/2013	Neurology	Dr. Stanley				

Fig. 4 Aggregated Data for the Problem of Epilepsy, Showing Lab and Medication Data and Links Which Give Access to Full Data for Imaging, Procedures and Notes

J Buchanan. Accelerating the benefits of the problem oriented medical record

Lab Test	LOINC Code	Medication	RxNorm C
GLUCOSE	2345-7	Acarbose	16681
GLUCOSE, FASTING	1558-6	Miglitol	30009
HEMOGLOBIN A1C	4548-4	Metformin HCl	235743
MICROALBUMIN/CREATININE RATIO	3000-4	Alogliptin Benzoate	1368000
CREATININE	2160-0	Linagliptin	1100699
E-GFR	33914-3	Saxagliptin HCl	1043560
POTASSIUM	2823-3	Sitagliptin Phosphate	621590
LDL CHOLESTEROL, CALCULATED	13457-7	Bromocriptine Mesylate	142426
LDL CHOLESTEROL	2089-1	Dulaglutide	1551291
LDL CHOLESTEROL, DIRECT	18262-6	Exenatide	60548
CHOLESTEROL	2093-3	Liraglutide	475968
HDL CHOLESTEROL	2085-9	Insulin Aspart	51428
CHOLESTEROL/HDL RATIO	2095-8	-	·
NON-HDL CHOLESTEROL	43396-1		
TRIGLYCERIDES	2571-8		

Fig. 5 Example of a Problem Concept Map

ASSESSMENT/PLAN:

Hypertension

Overall he has done pretty well in these last months. His weight has dropped substantially with diuresis from about 400 pounds last May to 365# last June and 362# at the end of September 2014. His weight in clinic today was 369#. Most notably, his echocardiogram has substantially improved. On September 24 his left ventricle appear normal in both size and function. There was mild concentric hypertrophy. His left atrium is quite dilated. Overall, this is a substantial improvement and reassuring that the hypertension may have been the primary cause of his LV dysfunction.

As of today, his regimen includes lisinopril 10 mg a day replacing the hydralazine and his bumetanide is down to 1 mg daily. His carvedilol is the same. His pressure in clinic today was low normal and we chose to reduce his amlodipine dose down to 5 mg daily.

Plan: Reduce amlodipine from 10 to 5 mg. No other changes for now.

Dyslipidemia


He is currently taking atorvastatin 40 mg. His most recent lipid panel has an LDL of 74, HDL 33 triglycerides 228. The biggest remaining issue here is his high triglyceride/low HDL which is being driven by likely a combination of his weight and his genetic background. At this point the bulk of his risk is addressed with the atorvastatin and we may no changes to that dose.

Plan: Focus on diet and exercise shifts to lose weight in the lipid number should follow in line. He met with our dietitian today as well at her notice added separately.

Renal insufficiency

Creatinine numbers were elevated on admission May 2014. He switched from hydralazine to lisinopril last spring and his numbers have remained moderately elevated from 1.74 (GFR 47) to 2.96 (GFR 25). Most recent is 1.73. He has been seen by the renal transplant team, and similar to his echocardiograms, they saw significant proteinuria last spring, but repeat urinary analysis in January showed no protein.

Plan: No specific changes recommended.



....

Dyslipidemia

He is currently taking atorvastatin 40 mg. His most recent lipid panel has an LDL of 74, HDL 33 triglycerides 228. The biggest remaining issue here is his high triglyceride/low HDL which is being driven by likely a combination of his weight and his genetic background. At this point the bulk of his risk is addressed with the atorvastatin and we may no changes to that dose.

Plan: Focus on diet and exercise shifts to lose weight in the lipid number should follow in line. He met with our dietitian today as well at her notice added separately.

Fig. 7 Natural Language Recognition of Problem Based Section Headings will Allow Storage, Retrieval and Display of the Assessment and Plan that are Particular to a Specific Problem

Item #	Link Type	Definition	Time of Creation	Storage Location	Example	Essential for Problem Concept Maps	Value
1.	Map Com- ponent Link of a Prob- lem Con- cept Map	A problem concept map component link is a con- nection specified in the system, between a cluster of SNOMED problem codes and codes for medi- cations, labs, imaging pro- cedure and diagnostic pro- cedures	Time of System Setup	System Level	COPD and Alpha- 1-Anti-Tryp- sin Level	Yes	Automatic; Does not depend on indication links and manual links which often do not exist
2.	Indication Link	An indication link is a con- nection between a medi- cation, treatment or other orderable item and a par- ticular visit diagnosis or problem for a patient. The link is created at the time of ordering.	Time of Or- dering	Patient Level	Albuterol or- dered with a specified in- dication of COPD	No	Reflects the in- tent of the pre- scribing provider
3.	Manual Link	A manual link is a connec- tion between a medi- cation, lab result, imaging result, diagnostic pro- cedure result or segment of narrative text and a particular problem for a patient.	Sometime After Or- dering or While Document- ing	Patient Level	Link be- tween COPD and a nar- rative prog- ress note about the COPD	No	Allows creation of a link to data or documen- tation in the ab- sence of a prob- lem concept map for that type of information

 Table 1
 Link Types Used for Dynamic Creation of Problem Oriented View

Item #	Type of Data	Map, Link or Function	Details
1.	Ancillary Data	Problem Concept Map (with component links)	System allows storage of <i>hard concept maps</i> that specify which ancillary data should be displayed when drilling down on particular problem (e.g. HgbA1C, creatinine, GFR and microalbumin/creatinine ratio for the problem of diabetes)
2.		View	System uses a dynamic display to view ancillary data (e.g. lab data, imaging reports, diagnostic procedure reports) in a problem oriented structure upon drilling down on a particular problem. Items are included in the display based on a <i>concept map relationship</i> .
3.	Medications and Other Treatments	Indication Link	Requirement for providers to specify an indication/prob- lem for all medications and other treatments, so they can be displayed when that problem is selected.
4.		Problem Concept Map (with component links)	System allows storage of <i>hard concept maps</i> that specify which medications should be displayed for a particular problem (e.g. for diabetes, this would include all insulins, sulfonylureas, biguanides, DPP-IV inhibitors, alpha-glu- cosidase inhibitors, thiazolidinediones, peptide analogs and glycosuric drugs). These medication maps are necessary because it will be some time before software can be implemented that requires an indication for every medication prescription.
5.		View	System uses a dynamic display to view medications and other treatments related to a given problem, upon drill- ing down on a particular problem. Items are included in the display based on an <i>indication link</i> or a <i>concept map</i> <i>relationship</i> .
6.	Documentation	Manual Link to Entire Note or to Assess- ment/Plan	Ability for documenting providers to use labels, links or templates to tie their documentation to a given prob- lem.
7.		Problem Concept Map (facilitates display of entire note)	When the user clicks a problem hyperlink, the system dy- namically generates a linked index of clinic notes, dis- charge summaries and consult notes related to that problem, utilizing the encounter diagnosis, provider specialty or clinic specialty for the individual piece of documentation.
8.		System Creates Tags for Assessment/Plan Sections	The system has the ability to utilize basic NLP to tag as- sessment/plan paragraphs with the name of existing pa- tient problems. When the user clicks a problem hyper- link, the system dynamically generates a linked index of these assessment/plan paragraphs.
9.		View	Ability to display a hyperlinked index of provider docu- mentation (clinic notes and hospital admission notes, daily progress notes and discharge summaries and oper- ative notes) in a problem oriented structure. Items popu- late the index based on <i>manual links</i> , <i>concept map rela-</i> <i>tionships</i> , or <i>tagged assessment/plan text</i> .

Table 2 Maps, Links and Functionality Needed for Generation of Problem Oriented View

References

- 1. Weed LL. Medical records that guide and teach. N Engl J Med 1968; 278: 593-600.
- 2. Weed LL. Medical records, medical education and Patient care: Press of the Case Western Reserve University, 1969.
- 3. Simons SM, Cillessen FH, Hazelzet JA. Determinants of a successful problem list to support the implementation of the problem-oriented medical record according to recent literature. BMC Medical Informatics and Decision Making 2016; 16(1): 1.
- 4. https://www.cms.gov/Regulations-and-Guidance/Legislation/EHRIncentivePrograms/downloads/ 3_Maintain_Problem_ListEP.pdf
- 5. Wright A, Maloney FL, Feblowitz JC. Clinician attitudes toward and use of electronic problem lists: a thematic analysis. BMC Medical informatics and Decision Making 2011: 1.
- 6. https://en.wikipedia.org/wiki/Network_effect
- Wright A, Pang J, Feblowitz JC, Maloney FL, Wilcox AR, McLoughlin KS, Ramelson H, Schneider L, Bates DW. Improving completeness of electronic problem lists through clinical decision support: a randomized, controlled trial. J Am Med Inform Assoc 2012; 19(4): 555-561.
- Wright A, Pang J, Feblowitz JC, Maloney FL, Wilcox AR, Ramelson HZ, Schneider LI, Bates DW. A method and knowledge base for automated inference of patient problems from structured data in an electronic medical record. J Am Med Inform Assoc 2011; 18(6): 859-867.
- 9. Wright A, Chen ES, Maloney FL. An automated technique for identifying associations between medications, laboratory results and problems. Journal of biomedical informatics 2010; 43 (6): 891–901.
- 10. Salmon P, Rappaport A, Bainbridge M, Hayes G, Williams J. Taking the problem oriented medical record forward. Proceedings of the AMIA Annual Fall Symposium 1996 (p. 463). American Medical Informatics Association.
- 11.Bainbridge M, Salmon P, Rappaport A, Hayes G, Williams J, Teasdale S. The Problem Oriented Medical Record-just a little more structure to help the world go round. Proceedings of the 1996 Annual Conference of PHCSG 1996 Sep.
- 12. Kaplan, DM. Clear writing, clear thinking and the disappearing art of the problem list. J Hosp Med 2007; 2(4): 199.
- 13. Kaplan, DM. Perspective: whither the problem list? Organ-based documentation and deficient synthesis by medical trainees. Academic Medicine 2010; 85(10): 1578–1582.
- 14. Don't Pave the Cowpath. http://www.nearfuture.io/dont-pave-the-cowpath/
- 15. Horsky J, Schiff GD, Johnston D, Mercincavage L, Bell D, Middleton B. Interface design principles for usable decision support: a targeted review of best practices for clinical prescribing interventions. Journal of Biomedical Informatics 2012; 45: 1202–1216.
- 16.Zeng Q, Cimino JJ. Evaluation of a system to identify relevant patient information and its impact on clinical information retrieval. Proc AMIA Annu Symp 1999: 642–646.
- 17.Zeng Q, Cimino JJ, Zou KH. Providing concept-oriented views for clinical data using a knowledge-based system: an evaluation. J Am Med Inform Assoc 2002; 9(3): 294–305.
- 18. Schiff GD, Seoane-Vazquez E, Wright A. Incorporating Indications into Medication Ordering--Time to Enter the Age of Reason. N Engl J Med 2016; 375(4): 306.
- 19. McDonald CJ, Huff SM, Suico JG, Hill G, Leavelle D, Aller R, Forrey A, Mercer K, DeMoor G, Hook J, Williams W. LOINC, a universal standard for identifying laboratory observations: a 5-year update. Clinical chemistry 2003; 49(4): 624-633.
- 20. What is IHTSDO? https://ihtsdo.freshdesk.com/support/solutions/articles/4000009519-what-is-ihtsdo.
- 21. Shaprio C and Varian HR. Information Rules. Harvard Business School Press. 1999.